



MILK

Nutritious by nature

The science behind
the health and nutritional impact
of milk and dairy foods




“Milk, Nutritious by Nature” is a European information initiative addressing nutrition and health issues.

The initiative’s objective is to disseminate relevant information and create a dialogue with key stakeholders such as high level scientists, health professionals, policy makers and journalists in relation to the nutrient richness and health impact of milk and dairy products and how those nutrients and other components come together in the dairy matrix to exert their beneficial effects. The aim is to build a clearer understanding of the role of milk and dairy products in a healthy, balanced diet across Europe.

This brochure for nutrition and health professionals is the result of collaboration between the nutritionists from the EMF member organisations to review the latest science on the role of milk and dairy foods in nutrition and health.

EMF comprises dairy organisations in eight countries - Austria, Belgium, Denmark, France, Ireland, Northern Ireland (UK), Norway and The Netherlands.



Milk, Nutritious by nature

Contents



Nutrient richness > 07

- 07 A unique package of essential nutrients
- 08 Contributing to diet quality in Europe
- 10 An important part of European dietary guidelines



Dairy and health > 12

- 13 Dairy matrix effects
- 15 Bone health
- 19 Blood pressure
- 22 Weight control
- 25 Type 2 diabetes
- 28 Cardiovascular disease
- 32 Cancer
- 35 Muscle mass maintenance in older people
- 38 Recovery after exercise



Conclusion > 40



References > 42

Milk, Nutritious by Nature

Milk and dairy foods have been an important part of the European diet and food culture for generations. They are naturally rich sources of a wide range of nutrients and make a significant contribution to nutrient intakes and diet quality.

Milk and dairy foods have also been linked to a variety of potential benefits in areas such as bone health, blood pressure and weight control, as well as type 2 diabetes, cardiovascular disease and colorectal cancer. They also have a role in sports nutrition and in helping to maintain muscle mass and function in older people.

This review summarises the scientific evidence of what makes Milk, Nutritious by Nature.



Nutrient richness

A unique package of essential nutrients

Milk and dairy foods are naturally rich sources of a wide range of essential nutrients. Many people associate milk and dairy with calcium and bone building but dairy foods offer much more extensive nutritional benefits.


Milk contains high quality protein, carbohydrate in the form of lactose, as well as many different fatty acids and a wide range of micronutrients including vitamins, minerals and trace elements.

Milk is a natural source of calcium, vitamin B12, riboflavin (vitamin B2), phosphorus and potassium.

It also contains smaller amounts of other nutrients including vitamin A, niacin, folate, vitamin B6, vitamin D, magnesium, selenium and zinc. In some, but not all, European countries milk is also a good source of iodine. The variation in iodine content is mainly due to differences in cows' diets between countries. The cows' diet can also affect the content of other nutrients, for example fatty acids and selenium.

Products made from milk such as yogurt, fermented milks and cheese also contain many of the nutrients which are present in milk. Many yogurts and cheeses, for example, are natural sources of protein, calcium, phosphorus and vitamins B2 and B12. Hard cheeses, in addition, also have zinc and vitamin A.

The nutrients present in milk and dairy foods are involved in a number of important functions in the body! These include the well-known benefits of calcium for bones and teeth, and protein for muscle. But dairy nutrients also play a part in nerve and muscle function, energy release, vision, blood clotting and red blood cell formation, digestion, blood pressure, skin health, the immune system, psychological function and in children's growth.



Contributing to diet quality in Europe

Given their rich nutrient content, it is not surprising that milk and dairy foods make an important contribution to the nutritional quality of the European diet. In many countries they are the main providers of calcium, contributing to 40% – 70% of the calcium intake in the European diet. Dairy also makes significant contributions to the intake of many other nutrients including high-quality protein, riboflavin, vitamin B12, phosphorus, vitamin A, iodine, zinc and potassium. For some countries and age groups, milk and dairy products also make worthwhile contributions to intakes of selenium, magnesium, niacin, folate and vitamin D.

Contribution (%) of dairy foods to nutrient intakes in adults in eight European countries

| Nutrients | Austria | Belgium | Denmark | France | Ireland | Netherlands | | Norway | UK |
|---------------|-------------|------------|------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | 18-65 years | 3-64 years | Total population | 18-79 years | 18-90 years | 31-50 years | 51-69 years | 18-70 years | 19-64 years |
| Protein | - | 19 | 24 | 16 | 13 | 23 | 24 | 22 | 14 |
| Calcium | 53 | 49 | 59 | 40 | 39 | 58 | 62 | 67 | 37 |
| Phosphorus | - | - | 33 | 23 | - | 32 | 34 | - | 22 |
| Potassium | - | - | 17 | 10 | - | 16 | 15 | 17 | 11 |
| Iodine | - | 16 | 30 | 21 | 44 | 16 | 16 | 60 | 35 |
| Zinc | - | - | 22 | 16 | 13 | 24 | 25 | - | 16 |
| Selenium | - | - | 16 | 7 | - | 14 | 14 | - | 6 |
| Magnesium | - | - | 14 | 9 | 11 | 14 | 15 | 14 | 10 |
| Vitamin A | - | - | 11 | 24 | 37 | 21 | 20 | 30 | 16 |
| Vitamin B2 | - | 32 | 41 | 26 | 29 | 42 | 42 | 37 | 29 |
| Vitamin B6 | - | 8 | 12 | 7 | | 12 | 12 | 11 | 8 |
| Vitamin B12 | - | 31 | 30 | 17 | 35 | 40 | 40 | 25 | 33 |
| Folate | - | 10 | 14 | 12 | 11 | 12 | 12 | 12 | 8 |
| Niacin | - | - | 17 | 2 | - | - | - | - | 7 |
| Vitamin D | - | 20 | 9 | 27 | 11 | 6 | 6 | 16 | 6 |
| Total fat | 14 | 18 | 17 | 17 | 13 | 18 | 20 | 26 | 14 |
| Saturated fat | 23 | 29 | 28 | 26 | 20 | 31 | 33 | 42 | 22 |
| Calories | - | 13 | 13 | 11 | 9 | 14 | 16 | 18 | 9 |

See page 42 for details of the sources of dietary data for individual countries.

The total amount of dairy foods consumed and the relative contribution of milk and other dairy products differs between countries, and, therefore, affects the contribution to nutrient intakes - for example, cheese makes a more important contribution to nutrient intake in France and Belgium than it does in the UK or Ireland where milk contributes proportionately more. Similarly, both the types and amounts of dairy consumed vary with age and so, therefore, do contributions to nutrient intakes. In general, milk contributes proportionally more to the diets of young children than to adults^{2,3}.

Some of the nutrients provided by dairy are not easily replaced by other foods without reducing the overall nutritional quality of the diet. For example, modelling dietary patterns to remove dairy and replace with non-dairy substitutes for calcium, resulted in lower amounts of several nutrients including protein, phosphorus, riboflavin, zinc and vitamin B12⁴.

Inadequate intakes of certain nutrients in the European diet are partly a consequence of low dairy consumption. A study providing information about micronutrient intakes across Europe shows that in some countries including France, the UK and, to a lesser extent, Denmark, intake of some micronutrients is inadequate, particularly among adolescents and young women⁵. These include calcium, iron, potassium, iodine, magnesium, selenium, zinc and riboflavin. Low intakes of magnesium, potassium, selenium were also evident in people over 60. With the exception of iron, these nutrients are all found in dairy foods, and the inadequate intakes are likely to be partly a reflection of low dairy intakes. In relation to teenagers and young women, for example, in many countries, milk drinking tends to decline in the teenage years especially in girls, with adverse consequences for intake of 'dairy' nutrients, which can be particularly important at this life-stage⁶.

The wide range of milk and dairy foods can suit different dietary needs. People with low energy requirements or those who are restricting their energy intake in order to lose weight, still need adequate amounts of protein, vitamins and minerals which low-fat dairy foods can provide. Others such as young children and the frail elderly may, in addition to their nutrient requirements, need to pay special attention to ensure their energy needs are met despite small or reduced appetites, and in this case whole milk and whole milk dairy products can be the most appropriate.

Nutrient density

It is increasingly recognised that a food's total nutrient content is important, rather than just the particular individual nutrients it contains. For example, in addition to specifying the quantities of nutrients needed in the diet, the latest Nordic Nutrition Recommendations emphasise the need to consider the particular foods that nutrients are obtained from and the quality of the food⁷.

One approach which takes into account the overall nutritional package of a food is nutrient density. The nutrient density of a food is usually defined as the ratio of essential nutrients, compared with the energy (calories) provided by the food – in some definitions these are also corrected for the presence of nutrients 'to limit' such as fat and sodium. The contributions of milk and milk products, particularly low-fat dairy, to nutrient intakes are relatively high compared to their calorie and fat contribution, giving them a favourable nutrient density score⁸.

Some scoring systems (such as the US Nutrient Rich Foods Index), which rank foods on the basis of their nutrient content per calorie, also allow the calculation of nutrients in relation to the cost of the food⁹. This is intended to help people choose diets that are both affordable and nutrient dense. Using such scoring systems, dairy foods, especially milk, have been shown to provide good nutritional value in relation to both calories and cost. This has been shown in European studies, as well as in the US¹⁰⁻¹².

An important part of European dietary guidelines

The milk and dairy food group forms an important part of food-based dietary guidelines across Europe, and dairy is recognised as one of the components of a healthy dietary pattern.

Food-based dietary guidelines

Dairy forms part of food-based dietary guidelines throughout Europe. The specific recommendations vary between countries, but on average 2 to 3 servings of milk and dairy foods a day are recommended for adults; often more for children and adolescents (around 3 to 4 servings), and in some cases more for pregnant women and older people too.

In some countries, dairy consumption falls short of the recommendations. In Belgium, for example, only 2% of the population (aged 3 to 64 years) meets the recommended intake of milk and milk products (except for cheese)¹³. In Ireland, adults (18 to 64 years) consume on average just over two portions from the 'milk, yogurt and cheese' food group a day compared to the recommended three servings; only 13% of adults meet this target¹⁴. In older Irish adults (60 years plus) the figure is lower still, at 3.5%¹⁵. Moreover, in many countries, milk and dairy consumption is declining. In France, for example, data from the CCAF Survey (French eating behaviours and consumption) reported a decrease in milk intake between 2013 and 2016 of 10% in children (3 to 17 years) and 18% in adults¹⁶.

Dietary patterns

Milk and dairy foods are recognised as part of a healthy dietary pattern. Increasingly, research is focusing on the effects of the whole diet on health and trying to distinguish the most favourable dietary patterns and habits. This is beginning to be translated into guidance for the public, including at national level. The 2012 Nordic Nutrition Recommendations, for example, have a stronger emphasis than previously on the whole diet, and describe a healthy dietary pattern as including plenty of vegetables, fruits and berries, pulses, regular intake of fish, vegetable oils, wholegrains, low-fat versions of dairy and meat, and limited intake of red and processed meat, sugar, salt and alcohol¹¹. Similarly, in the 2015 Dutch Dietary Guidelines, the Health Council of the Netherlands sets out what constitutes a healthy dietary pattern for the population, including milk and dairy foods¹⁷.





Dairy and health

Health impact of milk and dairy foods

Milk and dairy foods have been linked to a number of potential health benefits in areas such as blood pressure, bone health and weight control, as well as type 2 diabetes, cardiovascular disease and colorectal cancer. They also have a role in sports nutrition and in helping to maintain muscle mass and function in older people.


Ideally, the relationship between dairy foods and health would be measured by very large, very long-term intervention trials. In practice, however, the best available data are often from large, long-term observational studies. While not able to prove a cause and effect relationship, such prospective cohort studies and other types of observational data provide a 'real life' picture of a food in the diet and in a particular dietary pattern.

A number of analyses have pooled data from individual intervention or observational studies and the combined results increase statistical power and add weight to the results of single studies. Such meta-analyses provide the most robust evidence of the relationship between milk and dairy foods and health and, along with the recent reviews (both systematic and narrative), are discussed below. However, some individual studies conducted in European populations are also highlighted. These give an indication of the potential health effects of milk and dairy foods in relation to the particular amounts, types and patterns of consumption in Europe.

Intakes in European countries may be different to other parts of the world such as the United States or Asia where dairy consumption is often lower; they can also differ between European countries. In addition, the composition of milk and dairy products can vary between Europe and elsewhere depending, for instance, on animal feeding practices e.g. whether the cows are fed predominately on grain or fresh pasture. Fortification practices can also differ - e.g. vitamin D is routinely added to milk in the USA, and it is mandatory to do so in Canada. This is the case for low-fat milk in some Scandinavian countries including Norway but, on the whole, consumption of vitamin D-enriched milk is less commonplace in other parts of Europe. All of these factors may influence the results of research on the impact of dairy on health, and help explain differences between Europe and other parts of the world, and also within Europe.

Dairy matrix effects

It is increasingly recognised that the effects of milk and dairy foods on health extends beyond the individual nutrients they contain. Rather, the unique combination of nutrients and bioactive factors, and how they interact with each other in the dairy matrix, combine to produce the overall effect on health.



The food matrix

The ‘food matrix’ describes a food in terms of both its structure and its nutrient content, and how these interact together. Foods consist of a large number of different nutrients that are contained in a complex physical structure. The nature of the physical structure together with the mix of nutrients and bioactives can impact nutrient digestion, absorption and metabolism, affecting the overall nutritional and health properties of the food¹.

Whole foods or single nutrients

Nutrition research has traditionally focused on identifying the specific mechanisms and health impact of single nutrients – for example, saturated fatty acids in relation to blood lipids and cardiovascular disease (CVD). It has often then followed that nutrition policy is based on such associations - for example, recommendations to limit foods containing saturated fatty acids in order to reduce CVD risk.

More recently, however, nutrition research has shifted focus to examine the relationship of whole foods with health, including dairy foods (in both observational studies

and intervention studies as discussed above). This is based on the premise that we do not eat nutrients in isolation but as foods, and meals, and part of dietary patterns. From this research, a different picture has emerged in some cases than might be predicted from the nutrient content of the foods investigated. Cheese is a good example: despite its saturated fat (and salt) content, the majority of epidemiological studies report that cheese consumption does not increase the risk of CVD and may, in fact, be beneficial². Researchers have characterised this as a ‘food matrix’ effect¹. This recognises that the health effects of a food are much more complex than that of a single nutrient it contains or even a few nutrients. Rather, they are a function of both a food’s structure and its nutrient composition, and how these interact with each other.

Complex dairy matrices

Milk and dairy products are complex foods containing numerous nutrients and bioactive components. The rich dairy matrix of nutrients includes **protein, fat, lactose, calcium, phosphorus, potassium, vitamin B12, riboflavin (vitamin B2)** and many other vitamins and minerals. Dairy is also rich in bioactive

components - for example, small peptides with biological effects are produced when milk protein is digested in the intestine or during the fermentation process in foods such as cheese and yogurt³. Fermented dairy products also contain bacteria with the potential to produce beneficial short-chain fatty acids (SCFAs) in the gut⁴.

The nutritional and functional complexity in dairy food matrices is exemplified by milk fat, which has over 400 different fatty acids with different physiological properties^{3,5}. Components of the membrane which encloses the fat droplets in milk (the milk fat globule membrane; MFGM) also have functional effects, for example in relation to lipid metabolism⁶.

The matrix of different dairy foods will differ in the amounts and combinations of nutrients, including proportions of whey and casein proteins, fat and bioactives. Within a dairy food category, production methods and processes will also alter the composition of the matrix - for example, the length of ripening time for cheese affects the extent of protein breakdown and production of bioactive peptides¹.

In addition to their nutritional matrices, dairy foods also have complex physical matrices: from the solid matrix of cheese, gel-like structure of yogurt to liquid milk. Again, these differ within dairy categories - for instance, methods of production and ripening will influence the structure of the many different types of cheese.

The structure of a food matrix can have an impact on factors such as nutrient absorption and digestion and,

therefore, the metabolic response after eating. For example, rates of gastric emptying can differ for liquid compared to solid or semi-solid structures and so affect satiety and appetite responses⁷. The matrix structure can also influence protein absorption and digestion, for example, between casein contained in the milk matrix and given as a supplement⁸.


More than the sum of its nutrients
Given the complex nature of dairy matrices, it is perhaps not surprising that there is increasing evidence that the health effects of dairy foods extend beyond their constituent parts. Potential matrix effects are discussed in detail below. However, evidence to date suggests that milk and dairy matrices have specific beneficial effects on cardiometabolic disease risk, body weight and bone health¹.

For example, the effects of milk and dairy foods on bone health may be due in part to positive interactions of calcium, protein and phosphorus with each other and with lactose and bioactive peptides in the dairy matrices, rather than simply a 'calcium effect' as is often assumed. Similarly, the blood pressure lowering effect of milk may be the result of interactions between calcium, potassium, phosphorus and bioactive dairy peptides in the milk matrix. In relation to cheese, the explanation for the potential beneficial rather than harmful effects on CVD may again lie in interactions of the components of the cheese matrix including calcium, phosphorus, the milk fat globule membrane and starter cultures, which modify saturated fatty acid-induced increases in blood lipids.

More research is underway to investigate the health effects of the dairy matrix and to unravel the mechanisms and pathways through which the different components work together. However, the matrix concept underlines the importance of considering the health effects of milk and dairy as whole foods, alongside the individual components they contain. This is particularly important in relation to public health policy, and there is a growing recognition that dietary guidance should be based on evaluation of the health impact of whole foods, including dairy, rather than on single nutrients.

Bone health

The role of calcium in building and maintaining healthy bones is well established, and dairy foods are recognised as important sources of calcium, supplying up to two thirds of intake in the European diet. However, milk and dairy foods also contain other nutrients needed for bone health including protein, phosphorus and potassium (and vitamin D in the case of fortified dairy). Increasingly, the science indicates that nutrients in the dairy matrix work together to help maintain healthy bones. For example, there is some evidence that milk's calcium may offer longer lasting skeletal benefits than supplements due to its favourable calcium-phosphorus ratio, and that the calcium and protein in dairy have positive interactions on bone health.



Dairy and bone health in children and adolescents

Both observational and intervention studies provide evidence linking dairy consumption with bone health, especially in children and adolescents¹⁻³. In

children, a meta-analysis of studies investigating the effect of dairy products and dietary calcium (predominantly from dairy) on bone mineral content (BMC) reported that total body and lumbar spine BMC were significantly greater in children with higher intakes⁴. Dairy as part of an overall healthy dietary pattern has also been associated with beneficial effects on bone development⁵. Intervention studies which have specifically used milk or dairy foods are limited compared with those for calcium supplements, but positive effects have been reported; including in French, Finnish and British children⁶⁻⁸. For example, significant improvements in bone mineral acquisition compared to control subjects were observed in adolescent British girls who were given 568ml (one pint) of milk a day for 18 months⁹. The corollary is that milk avoidance in children has been associated with increased risk of fracture and poorer bone health^{10,11}. In a 2016 position statement on lifestyle choices that promote maximal bone health from childhood through adolescence, the National Osteoporosis Foundation concludes that there is 'good evidence' for a role for dairy consumption (the 'best evidence' is for the positive effects of calcium intake and physical activity)¹². A number of retrospective studies, although not all, have found that milk consumption in childhood and adolescence is related to better bone health and / or reduced risk of fracture later in life¹³. There are limitations to such studies, however, including accurate recall of childhood milk and dairy intake.

Dairy and bone health in adults

For adults, the majority of observational studies report either a positive association between milk and milk products and BMC or bone mineral density (BMD) or a neutral outcome¹². Again, randomised

controlled trials using milk and dairy foods are limited compared with calcium supplementation ones, and longer-term trials and meta-analyses are needed. In such studies the outcome will depend on factors such as the age of the subject, relation to the menopause for women, initial dairy intakes and so on¹³. Intervention studies have, however, reported positive associations between increase in dairy food intake and BMC or BMD, and reductions in bone turnover markers^{14,15}. On the whole, there is support for favourable effects of dairy on measures of bone health in adults³.

Fracture risk

With respect to fracture risk, the effects of dairy are less clear. This may be due to heterogeneity in the study

designs, duration, participants' age, and other confounding factors such as vitamin D status and physical activity. A meta-analysis of prospective cohort studies published in 2011, found no overall association between adult milk intake and hip fracture risk in women; insufficient data was available in men¹⁶. However, the data on women were disproportionately influenced by one study from Sweden; when the authors excluded this study from the analysis there was a marginally significant 5% lower hip fracture risk for each glass of milk consumed per day. A subsequent publication in 2014 utilising data from the same Swedish cohort of 61,000 women but with a longer follow up (around 20 years) found fermented milk products (yogurt and other soured milk products) and cheese consumption were associated with a significant decrease in fracture incidence¹⁷. However, high intakes of milk (three or more glasses/day; more than 600 ml/day) were associated with increased fracture rate. It is worth noting that when the dietary questionnaires were performed (1987–90 and 1997) milk in Sweden was fortified with high dose of

vitamin A: such levels of vitamin A intake have been linked to an increased risk of fracture. A study of Finnish women has reported that milk avoidance (because of lactose intolerance) was associated with increased fracture risk¹⁸.

No dairy intervention trials on fracture risk are available because of the feasibility of carrying out such a study; however, calcium supplementation trials do exist.

A meta-analysis of 17 randomised trials concluded that calcium supplementation (with or without vitamin D) decreases fracture risk by 12% in people aged 50 years and older¹⁹. The fracture risk reduction was greater (24%) in trials when compliance was high; also in participants older than 70 years and whose daily calcium intake was initially low.

Potential dairy matrix mechanisms

The importance of calcium in bone development and maintenance is well established^{1,13}.

Milk and milk products make the largest contribution to calcium intake in the European diet. Few other foods naturally contain as much calcium, and dairy sources are some of the most bioavailable²⁰. While it is sometimes assumed that supplementation with the same amount of calcium from different sources - e.g. milk, foods fortified with calcium and calcium supplements – have comparable effects on bone health, there is some evidence of a beneficial 'dairy matrix effect'²¹. Dairy foods may have greater benefits than the equivalent calcium in the form of supplements. In adolescent girls, for example, it has been estimated that bone mineral density increases by up to 10% when 700mg extra calcium is provided as dairy foods, compared with an increase of 1% to 5% when the same quantity of calcium is given as a supplement²². Similarly, using cheese to increase calcium intake in 10- to 12-year-old Finnish girls resulted in a greater increase in bone mineral density compared to either a calcium supplement or a calcium plus vitamin D supplement⁷. Another randomised controlled trial investigating the effect of dairy products providing 1,200mg calcium a day or an equivalent calcium supplement on

markers of bone metabolism and BMD found that after 12 months, the dairy intervention group had greater improvements in pelvis, spine and total BMD than the calcium supplement group¹⁴.

The greater benefits of dairy may be due to the presence of other nutrients in the dairy matrix which are important for bone health such as protein and phosphorus, and their interactions with calcium.

Protein is essential for bone development in children and is needed for the maintenance of normal bones in adults since amino acids are required for the synthesis of intracellular and extracellular bone proteins. Older adults consuming a protein-restricted diet are at higher risk for bone loss and fractures, and sufficient protein intake is recommended in guidelines for maintaining skeletal health²³⁻²⁶. There has been some controversy around the adverse effects of high protein intakes, but it is now established that although a high-protein diet increases urinary calcium excretion this does not result in a negative skeletal calcium balance, or bone loss²⁴⁻²⁸. Indeed, recent research suggests higher protein intakes are beneficial to bone health, especially when calcium intake is also adequate²⁹. For example, in the US Osteoporotic Fractures in Men study, greater intakes of dairy protein were associated with a decreased risk of hip fracture³⁰. Similarly, in healthy postmenopausal women, dairy protein intakes were positively associated with measures of bone strength and microstructure³¹. The potential mechanisms for this include protein enhancing calcium balance by stimulating intestinal calcium absorption, both directly and indirectly via an IGF1-vitamin D link. Part of the explanation of the greater effectiveness of dairy calcium versus supplements may also be due to better absorption of calcium because of the presence of lactose and / or casein phosphopeptides in the dairy matrix⁷. Fermented dairy products may additionally enhance calcium absorption through positive alterations in the gut microbiota³².

In addition to potentially larger effects, it has been suggested that the skeletal benefits of dairy calcium may persist longer than from calcium supplements²⁵.

In a study of 8-year-old French girls, the benefits of milk-extracted calcium phosphate on bone mass remained 3.5 years post supplementation⁴. This has not been the case after supplementation with calcium salts (such as citrate malate or carbonate). Part of the explanation may lie in the favourable calcium to phosphorus ratio in milk. Phosphorus

(as inorganic phosphate) is an important structural component of bone and an adequate intake is necessary for bone growth and development, and the maintenance of normal bones in later life²⁴. Although a high intake, if accompanied by low calcium (in a ratio of about 4:1) may be deleterious to bone, the phosphorus to calcium ratio of milk (0.8:1) can enhance calcium balance by stimulating renal tubular reabsorption of calcium and lead to positive effects on bone³³.

Other nutrients in milk and dairy foods are also involved in bone health including magnesium, potassium, vitamin K2 and zinc, as well as vitamin D in fortified dairy¹. Increasingly, the science indicates that the nutrients in the dairy matrix may work together to help maintain healthy bones. Simply in terms of the quantities of 'bone' nutrients in dairy, however, it has been suggested that it is difficult to devise a diet that is 'bone healthy' without including three servings of dairy a day¹.

Blood pressure

Observational and clinical studies suggest that milk and dairy intake, particularly low-fat dairy, could have a beneficial effect on blood pressure and contribute to the prevention of hypertension. The DASH (Dietary Approaches to Stop Hypertension) diet, which focuses on fruit and vegetables and low-fat dairy foods, has been found to be an effective way to lower blood pressure.

Milk and dairy foods contain several nutrients and other bioactive components in the dairy matrix including calcium, potassium, phosphorus and bioactive peptides, which may be involved, individually or in combination, in the beneficial effects on blood pressure.



Observational studies

A number of observational studies have noted an association between milk and dairy intake and lower blood pressure, including in European populations.

For example, in Welsh men, milk intake predicted systolic blood pressure: in the group with the highest milk intake (around a pint / 586ml of whole milk per day), systolic blood pressure was 10.4 mmHg lower than those who drank little or no milk after a 23-year follow-up¹. Similarly, for older people (over 55 years) in the Rotterdam Study, the risk of hypertension decreased with increasing low-fat dairy consumption; although no relationship between dairy and blood pressure was seen in a wider age range of the Dutch population (20–65 years)^{2,3}. In the French DESIR cohort, dairy (milk or yogurt) and cheese consumption were associated with lower diastolic blood pressure after the 9-year follow-up⁴. Cross-sectional data from the French cohort of the MONICA study also showed higher dairy intake was associated with lower systolic blood pressure⁵. A recent analysis of the National Adult Nutrition Survey in Ireland also reports an association between higher total dairy, and specifically milk, intake and lower systolic and diastolic blood pressure⁶.

In a meta-analysis of five cohort studies, consumption of dairy foods was associated with a 13% reduced risk of elevated blood pressure⁷.

Further analysis suggested that the effect may be driven by low-fat dairy and ‘fluid’ dairy (defined as milk and yogurt); cheese and full-fat dairy foods had no association with risk of high blood pressure. In another meta-analysis, of nine prospective cohort studies in 2012, dairy consumption was also associated with a reduced risk of hypertension⁸. Again, the effects were specific for low-fat dairy and milk (3% reduction per 200g/day) whereas there was no association for cheese, full-fat dairy, total fermented dairy and, in this case, for yogurt. Both meta-analyses suggest that milk and low-fat dairy could contribute to the prevention of hypertension. This concurred with the US Dietary Guidelines Advisory Committee’s assessment

of the science in 2010 which concluded that there was a moderate body of evidence that high intake of milk and milk products is associated with lower blood pressure⁹; subsequent studies have strengthened this conclusion. A systematic review published in 2016 of the association between dairy product consumption and risk of various cardiovascular-related clinical outcomes reports favourable associations between intakes of total dairy, low-fat dairy and milk and the risk of hypertension¹⁰.

The DASH diet

The best known intervention involving dairy, the DASH (Dietary Approaches to Stop Hypertension) diet, has proved an effective way to lower blood pressure in those with and without hypertension^{11–13}.

The DASH eating plan, which emphasises fruit, vegetables, wholegrains and low-fat dairy products (around 3 servings a day), has been widely promoted in the USA for the prevention and treatment of high blood pressure¹⁴. Although this intervention was first conducted as a feeding trial, further studies have found that DASH dietary advice can also be effective at lowering blood pressure in free-living populations^{15,16}. European guidelines on the management of hypertension now also include a ‘DASH diet’ approach recommending increased consumption of vegetables, fruits, and low-fat dairy products¹⁷. Observational studies have suggested that in children too, a ‘DASH’ dietary pattern may have beneficial effects on blood pressure^{18,19}. A recent trial has also indicated that when higher-fat dairy foods are incorporated into the DASH pattern, blood pressure lowering effects are still evident²⁰. In addition, three controlled trials which examined the impact of dairy products per se on blood pressure (rather than as part of a dietary pattern such as DASH) reported beneficial effects^{21–23}. These included an intervention using full-fat hard cheese (two months of 30g/day of Grana Padano) compared with a placebo consisting of flavoured bread mixed with fats and salts in the same concentrations as the cheese²³.

Potential dairy matrix mechanisms

Milk and dairy foods contain several nutrients and other bioactive components which have been associated with blood pressure control. Recent

research has focused on the importance of **bioactive peptides** in the regulation of blood pressure, including those from dairy. For example, a group of peptides (lactotriptides), released from milk and dairy products during digestion of casein proteins in the gut or by fermentation, have been shown to have anti-hypertensive properties and to regulate blood pressure by inhibiting ACE-1, a potent vasoconstrictor²⁴⁻²⁶. Similarly, in a recent clinical trial, whey protein also lowered blood pressure and improved endothelial function in adults with pre-hypertension and mild hypertension²⁷.

Blood pressure-lowering effects of the B vitamin riboflavin, of which milk is a good source, have also been reported²⁸.

This reflects the role of riboflavin in regulation of homocysteine levels in those with a genetic defect in homocysteine metabolism (about 10% of the European population); an elevated level of homocysteine has been associated with hypertension²⁹.

The minerals in milk including calcium, potassium and magnesium are also linked to blood pressure regulation

through their effects on intracellular mechanisms and production of vasodilators^{30,31}. **Calcium**, for example, may have a direct impact on blood pressure through effects on vascular smooth muscle, as well as through parathyroid hormone (PTH) and vitamin D secretion, and increased sodium excretion³¹. **Phosphorus** in the dairy matrix may be involved too³¹. It has been reported that phosphorus from dairy products, but not from other sources, is associated with lower baseline blood pressure and reduced risk of hypertension³². This may indicate that the benefits of phosphorus are dependent on interactions with other dairy components. Indeed, it is likely that the blood pressure lowering effects of milk and dairy products are the results of interactions between the constituents of the dairy matrix³³.

The weight of the evidence to date suggests that milk and dairy foods, particularly low-fat dairy, can help lower blood pressure and contribute to the prevention of hypertension. This is important given that high blood pressure is a major risk factor for cardiovascular disease, particularly stroke, and even values at the high end of the normal range increase the risk. Around 30% to 45% of the European population has hypertension, with a steep increase with ageing, so even small reductions in prevalence could have public health benefit¹⁷.

Weight control

Contrary to the popular perception that dairy foods are ‘fattening’, a growing body of research suggests that milk and dairy foods may have a positive role in weight control in both adults and children. A number of observational studies have reported that dairy-rich diets are associated with lower body weight, less body fat, less abdominal obesity and less fat gain over time. Overall, results from intervention studies suggest that including dairy foods in a weight-reducing diet may enhance weight loss.

Dairy’s calcium and protein are likely to be involved in its effects on energy balance, including through influences on appetite and satiety, fat absorption and energy use. From the available evidence it seems likely that for those trying to lose weight, avoiding dairy or having a low dairy intake may be counterproductive.



Observational studies

Scientific evidence is accumulating to suggest that milk and dairy foods may have a positive role in weight control in both children and adults. A

relationship between calcium and weight was first noticed in a cross-sectional analysis more than 30 years ago¹. However, it was not until the late 1990s, when a possible mechanism linking calcium and weight was proposed, that research in this area gathered pace². Since then, a number of other cross-sectional studies have noted a similar negative relationship between calcium and/or dairy foods and weight, body fat and abdominal obesity including some in European populations³⁻¹⁰.

For example, in a study of early-postmenopausal Italian women, those in the highest quartile of dairy intake had a lower BMI than those who ate the least dairy foods⁸. Similarly, data from the ‘Observation of Cardiovascular Risk Factors in Luxembourg’ survey found dairy intake was associated with a reduced likelihood of both overall obesity and abdominal obesity⁹. A recent analysis of the National Adult Nutrition Survey in Ireland also reports that higher total dairy intake, as well as higher milk and yogurt intake, were associated with lower measures of body fatness, including waist-to-hip ratio, % body fat and waist circumference¹⁰.

Some prospective studies too suggest a modest protective effect of dairy product consumption on the amount of weight gained over time¹¹. For example, a Swedish study which examined the association between dairy intake and weight change in 19,000 peri-menopausal women over nine years, found that regularly eating one or more servings a day of cheese or whole / fermented milk (3% fat) was associated with lower weight gains in normal weight women¹². In overweight French men, milk and yogurt intake were related to lower gains in weight and waist circumference over six years¹³.

A narrative review in 2011, concluded that data from observational studies, although not completely consistent, were suggestive of a protective effect of dairy consumption on the risk of overweight and obesity in adults⁶.

A systematic review of prospective studies in the same year reached a similar conclusion with regard to dairy and the risk of weight gain¹¹. A more recent analysis also reports that dairy consumption was not adversely related to changes in body weight; yogurt, in particular, showed a beneficial effect, where higher intakes were associated with a reduced risk of obesity, changes in body weight and waist circumference¹⁴. Another systematic analysis, which assessed the relationship between the intake of dairy fat and high-fat dairy foods with obesity, found no evidence that dairy fat intake adversely affects weight or obesity risk¹⁵. In fact, in the majority of observational studies (11 out of 16 studies), a dietary pattern with a high-fat dairy intake was associated with lower weight and other measures of adiposity.

Studies in children and adolescents

For children and adolescents there are fewer observational studies available, but on the whole these too indicate either a beneficial or neutral effect of dairy consumption on body weight or body composition^{11,16}. In UK children, for example, higher dairy consumption during preadolescence (10 years old) was not

associated with excess fat gain during early adolescence (up to 13 years) and, in fact, appeared to have a protective effect¹⁷. A recent meta-analysis looking at the long-term association between dairy consumption and risk of childhood obesity concludes that dairy is protective¹⁸.

Intervention studies

Results from intervention studies strengthen the observational data. A meta-analysis of randomised control trials in 2012 found that inclusion of dairy products in calorie-restricted diets led to a significantly greater reduction in body weight, waist circumference, and fat mass, while maintaining lean body mass, compared with low-dairy weight-loss diets¹⁹. A more recent meta-analysis in 2016 reached the same conclusion: increased dairy intake as part of energy-restricted diets resulted in greater loss of body weight and fat while helping to reduce the amount of lean mass lost²⁰. Other meta-analyses also report dairy consumption, in the short-term, coupled with calorie-restriction, has a beneficial impact on body fat reduction^{21,22}. Increasing dairy intake without calorie-restriction had no effect on weight^{19,21,22}. It is an important point that, in contrast to the popular perception, extra milk and milk products did not result in weight gain, and, when coupled with calorie restriction, enhanced weight loss.

Again, there are fewer intervention studies in children and adolescents and no meta-analysis of results, however, the majority of the research indicates that the effect of milk and dairy intake on body weight and body composition in children and adolescents is beneficial or neutral¹⁶.

Potential dairy matrix mechanisms

Possible mechanisms to explain the effects of dairy on weight control initially centred on calcium and particularly the hypothesis that calcium may alter fat cell function and fat oxidation: stimulating lipolysis (fat breakdown), reducing lipogenesis (fat synthesis) and increasing fat oxidation^{2,23,24}. More recently it has been

suggested that calcium may also work by binding fat in the intestine and increasing its excretion from the body and so decreasing fat (and therefore calorie) absorption²⁵. Calcium in the dairy food matrix may be more effective in this respect than other forms of calcium²⁶. Calcium-driven effects on appetite have also been postulated: a low calcium intake may trigger hunger and impair weight loss on energy restricted diets²⁷. These potential effects of calcium are not mutually exclusive and multiple mechanisms may be involved.

Other components of the dairy matrix are also likely to be involved in dairy's beneficial effects²⁸. Studies have reported that the weight-loss effects of milk and yogurt, for example, are greater than for the equivalent calcium supplement^{2,29}. A probable candidate is dairy's


protein. Several investigations indicate a role of protein in weight loss and weight maintenance including the European DIOGENES study³⁰. Dairy protein may have positive effects on satiety, as well as benefits on 'muscle sparing' helping to maintain lean body mass during energy restriction^{6,31}. Branched-chain amino acids, of which dairy foods, and particularly whey protein, are rich sources may be important in this respect³¹. Given associations between dairy fat and lower body weight, it has also been suggested that some fatty acids, principally **medium-chain fatty acids**, may have anti-obesity effects, including through lipogenesis and satiety^{32,33}. Moreover, there is preliminary evidence that short-chain fatty acid production in the gut, which is favoured by fermented dairy foods such as cheese, may have positive effects on appetite regulation³⁴.

Further research will help to fully elucidate the relationship between dairy consumption and body weight including amounts and types of dairy foods and possible threshold effects. Nevertheless, from the available data, it seems likely that those trying to maintain a healthy weight, and particularly for those trying to lose weight, a low dairy intake is likely to be counterproductive. Indeed, recent guidelines on the 'Dietary Treatment of Obesity' from the Swedish Council on Health Technology Assessment, advise that a higher intake of dairy products (mainly milk) during energy restriction can lead to weight loss for both adults and children³⁵.

Type 2 diabetes

Evidence is building that dairy foods may help reduce the risk of type 2 diabetes. Protective associations have been reported for total dairy consumption as well as for both low-fat and regular-fat dairy, and are particularly pronounced for fermented products such as yogurt and cheese.

Several components of dairy foods could potentially be involved in the reduced risk including protein, calcium, magnesium and dairy fatty acids, along with mechanisms related to fermentation. These are not mutually exclusive and may well interact in the dairy matrix to produce the beneficial effects. Since the number of people with type 2 diabetes is rising in European populations, even a small protective effect of dairy as part of a healthy diet could have important public health implications.



Observational studies

A number of epidemiological studies have reported that intake of milk and dairy food is associated with a reduced risk of developing type 2 diabetes.

An overview of four prospective cohort studies in 2007 showed a reduction of about 15% in the risk of diabetes in the highest dairy consumers (3 to 5 compared with less than 1.5 servings/day)¹. Similar reductions were reported in overviews in 2010 and 2011 based on five and seven cohorts respectively^{2,3}. In the latter meta-analysis, the protective effect of dairy was largely attributable to low-fat dairy foods, and a further dose-response analysis (involving three of the studies) suggested a 10% reduction in type 2 diabetes for every additional serving of low-fat dairy a day.

Since these overviews, several more prospective studies looking at the relationship between dairy consumption and type 2 diabetes have been published, including for European populations. The EPIC-InterAct Study (an investigation in eight European countries to identify the genes and lifestyle factors that influence risk of type 2 diabetes) found that both cheese consumption and fermented dairy product intake (cheese, yogurt, and thick fermented milk combined) were associated with a reduced risk of diabetes⁴. Similarly, in another EPIC cohort, this time in the UK and using 7-day food diaries, low-fat fermented dairy intake, largely driven by yogurt, was associated with a decreased risk of type 2 diabetes: 24% less between the highest and lowest consumers⁵. The PREDIMED study in an elderly Mediterranean population at high cardiovascular risk also reports that a healthy dietary pattern incorporating dairy products and, again, particularly yogurt, was protective against type 2 diabetes⁶. In the French DESIR cohort (Data from the Epidemiological Study on the Insulin Resistance Syndrome), those who consumed more than three servings of milk or yogurt a day had a lower risk of type 2 diabetes compared with those who consumed less than one serving a day⁷. Two other recent studies, in Danish and British cohorts, found no relationship

between dairy intake and diabetes^{8,9}. The Danish data did suggest a beneficial effect of cheese and fermented dairy on glucose regulation measures but these did not translate into a significant association with type 2 diabetes⁸. Another Danish study looking specifically at milk intake found no relationship with diabetes risk¹⁰. In the Dutch Maastricht Study, a high intake of full-fat products was positively related to diabetes risk¹¹. However, individuals with a high total dairy consumption had lower risk of type 2 diabetes, and those with a high consumption of skimmed and fermented products had lower risk of impaired glucose metabolism¹¹.

A systematic review and dose-response meta-analyses published in 2013 utilised the available data to investigate in more detail the relation between the intake of individual dairy foods and diabetes risk, as well as between regular- and low-fat dairy¹². Seventeen cohort studies were included in the overall meta-analysis and a significant reduction in the risk of type 2 diabetes was reported with increasing intakes of total dairy products and low-fat dairy products (up to 300g - 400g per day) and with increasing intake of cheese (up to around 50g per day). Another meta-analysis published in 2013 also found that total dairy product consumption, low-fat dairy products and cheese were associated with a reduced risk of diabetes, and additionally in this overview, so was yogurt¹³. In this case, total dairy and low-fat dairy intake were associated with 6% and 12% respectively, lower risk of type 2 diabetes per 200g a day consumption, with most of the risk reduction occurring with intakes up to about 200g per day for total dairy, and 300g per day for low-fat dairy.

A more recent analysis of cohort studies to inform the 2015 Dutch Dietary Guidelines concluded that there is convincing evidence that yogurt consumption is associated with a lower risk of type 2 diabetes; yogurt intake of more than 60g a day was associated with a 15% reduction in diabetes risk, compared

with intakes less than 10g per day¹⁴. Similarly, a meta-analysis in 2016 which summarised the evidence from 22 prospective cohort studies found yogurt consumption was associated with a reduced risk of type 2 diabetes at 80g/day, but not with higher intakes¹⁵. In this analysis, total dairy food consumption was also associated with a lower risk, mainly attributable to low-fat dairy foods; there were no associations with either milk or cheese intake. The same conclusion was reached from a systematic review published in 2016; namely, that there is high-quality evidence to support favourable associations between yogurt and low-fat dairy intake and reduced risk of type 2 diabetes¹⁶.

Intervention studies

Randomised control trials are needed to further confirm the results of the cohort studies. One recent small-scale trial found that consumption of four servings of low-fat milk and yogurt a day for six months improved plasma insulin and insulin resistance without negatively impacting on body weight or blood lipids¹⁷. Similarly, a six-week trial of three servings of low-fat dairy a day resulted in improvements in fasting glucose in men¹⁸.

Potential dairy matrix mechanisms

Several components of the dairy matrix could potentially be involved in the protective relationship between dairy and type 2 diabetes. These are not mutually exclusive and a combination of mechanisms may well produce beneficial effects on glycaemic control

and other risk factors. These include dairy minerals such as **calcium and magnesium**: both have a role in regulating insulin-mediated intracellular processes^{1,19,20}. **Protein** in dairy has also been shown to have beneficial effects on insulin secretion and blood glucose control²¹⁻²⁴. The **bioactive peptides** that may be involved result from digestion of milk protein in the gut and from the fermentation process in foods such as cheese, yogurt and fermented milk. Similarly, a form of **vitamin K** (vitamin K2; part of the menaquinone family) which is associated with fermented dairy foods has been linked to reduced risk of type 2 diabetes²⁵.

In addition, dairy fatty acids may play a role. Certainly, evidence from cohort studies suggests that the positive effect of dairy on diabetes risk is not restricted to low-fat dairy foods but also includes higher fat ones such as cheese. Data from the Malmö Diet and Cancer cohort, for example, indicated a decreased risk of type 2 diabetes with high intakes of regular-fat dairy foods but not with low-fat dairy²⁶. A number of recent studies have found that the dairy-derived fatty acid trans-palmitoleate is associated with a substantially lower incidence of diabetes (60% less in one study)²⁷⁻²⁹. Other fatty acids in dairy have also been associated with reduce diabetes risk including odd-chain saturated fatty acids pentadecanoic acid (C15:0) and heptadecanoic acid (C17:0) and short-chain fatty acids (SCFA) including butyrate (C4:0); the branched-chain dairy fatty acid phytanic acid has been reported to have anti-diabetic effects including increasing insulin sensitivity³⁰⁻³².

Dairy may indirectly modify diabetes risk through beneficial effects on weight, body fat and central adiposity, and on muscle mass and function. Additional research will help to differentiate the effects of individual dairy foods and components of the dairy matrix for diabetes risk particularly the relevance of fermented dairy products such as yogurt. Randomised control trials to confirm the results of cohort studies would also be useful. However, the protective effect of dairy suggested by evidence to date, has important public health implications given that there are already around 60 million people with diabetes in the European Region, and these numbers are increasing³³.

Cardiovascular disease

The focus on dairy foods and cardiovascular disease (CVD) is often in relation to saturated fat. There is an assumption that because some dairy foods contain saturated fatty acids and dairy in general contributes to saturated fat intake in the diet, that it also increases the risk of cardiovascular disease. Yet the majority of epidemiological studies have shown no adverse effects of regularly consuming milk and dairy foods such as yogurt and cheese on cardiovascular health, irrespective of fat content. In fact, in some cases a cardio-protective effect has been observed.

The explanation for this may lie in the complex composition of milk and dairy foods which, in addition to saturated fat, contain other nutrients and bioactive components such as calcium, potassium and bioactive peptides in the dairy matrix which may be beneficial to cardiovascular health. In addition, the overall fatty acid profile of milk and dairy may not have the detrimental effect on blood lipids or other cardiovascular parameters that has been assumed.



Observational studies

In the absence of long-term intervention trials, the best available data on the relationship between dairy foods and cardiovascular disease are from large, long-term observational studies.

There are several studies of this type in European populations. For example, a prospective cohort study from the UK reports that men who drank the most milk (around a pint / 586ml of whole milk a day) had fewer heart attacks and fewer strokes than those who had little or no milk in their diets¹. The very large Netherlands Cohort Study, consisting of over 120,000 men and women, showed no association between total milk product consumption and stroke mortality, although butter and dairy fat was associated with a small (7%) increased risk of all-cause and heart disease mortality among women². Recent data from the smaller Dutch Hoorn Study also found that overall dairy intake was not associated with CVD mortality but the intake of high-fat dairy products was³. In contrast, the Rotterdam Study reports that in an older Dutch population, high-fat dairy was associated with a reduced risk of fatal stroke; total dairy consumption or the intake of specific dairy products was not related to CVD events⁴.

In another Dutch prospective cohort, there was again no evidence that dairy products increased risk of heart disease or stroke⁵. In fact, high intakes of total and low-fat dairy were associated with a lower coronary heart disease (CHD) risk in participants without hypertension. Low-fat dairy consumption was also associated with reduced risk of stroke in cohorts of Swedish men and women⁶. In the same cohorts, a high intake of fermented milk (yogurt and cultured sour milk) was found to reduce CVD risk⁷. Fermented milk and cheese were also associated with reduced cardiovascular disease mortality⁸. There was, however, an increased CVD mortality risk reported in this study in those drinking three or more glasses of (non-

fermented) milk a day compared with less than one glass. The reason for this discrepancy in the same cohort is not clear and the authors urge a cautious interpretation of the results. Moreover, when these data were re-examined, milk consumption was associated with a lower risk of CVD mortality⁹. In line with the majority of epidemiological studies, a recent Danish investigation reports no adverse effects of dairy on cardiovascular health¹⁰. The French MONICA project conducted over 14 years, found that dairy consumption (particularly milk intake) as part of a diverse, healthy diet was associated with the lowest mortality rate mostly due to reduced cardiovascular deaths¹¹.

Meta-analyses for milk

A number of analyses have pooled the data from individual prospective studies such as these and their results strengthen the evidence that regularly consuming milk and other dairy products does not increase risk of cardiovascular disease

and may even have a protective effect^{1,13-20}. In relation to milk, an overview conducted in 2010 concluded that milk drinking is not harmful and may be associated with a small but worthwhile reduction in risk of coronary heart disease (8%) and a more substantial reduction in stroke risk (21%) for those who drank the most milk compared with those who drank the least¹³. The pooled results of seventeen studies in 2011 also found milk intake was associated with a small potential reduction in overall cardiovascular risk of 6% for each 200ml of milk consumed a day¹⁴. This analysis found no association between high intakes of either regular-fat or low-fat dairy products and increased risk of death from cardiovascular disease. Similarly, systematic reviews in 2015 and 2017 examining milk consumption and cardiovascular disease mortality observed no consistent association^{15,16}. This was also the conclusion for milk and CVD risk in a meta-analysis published in 2016; milk intake was found to be neutral with respect to risk of stroke and coronary artery disease¹⁷.

Meta-analyses for dairy products

Meta-analyses also support neutral or beneficial effects of other dairy foods on cardiovascular disease.

Twenty two prospective cohort studies were included in an analysis published in 2015 which examined stroke and CHD incidence in relation to intake of individual dairy foods, and to low- and regular-fat dairy¹⁸. Cheese consumption was associated with a 16% decreased heart disease risk, and both cheese and low-fat dairy foods were associated with reduced risk of stroke (9% and 7% respectively). An earlier meta-analysis in 2014 looking specifically at stroke also reported similar reductions in risk with low-fat dairy (9%) and cheese intake (6%) and, in addition, with total dairy (12%) and fermented milk (20%)¹⁹. Similarly, in another meta-analysis including 18 studies which had examined dairy intake and stroke risk, milk and cheese consumption were associated with reduced risk of stroke; risk reductions were maximal around 125g/day for milk (16%) and from 25 g/day upwards for cheese (9%)²⁰. Cheese was also associated with a lower risk of stroke (13%) in a meta-analysis published in 2016, as was total dairy intake (9%)²¹. In addition, cheese intake was associated with an 18% reduced risk of coronary heart disease. A beneficial effect of cheese was also supported by a meta-analysis of prospective cohort studies published in 2017 in which cheese intake was associated with 10%, 14% and 10% reduced risks of total CVD, CHD and stroke respectively²². A recent systematic review of the association between dairy product consumption and risk of various cardiovascular-related clinical outcomes also reports favourable associations between intakes of total dairy, low-fat dairy, cheese, and fermented dairy and the risk of stroke¹⁷. Similarly, a dose-response meta-analysis in 2017 combining data from 29 prospective cohort studies demonstrated neutral associations between dairy products and cardiovascular mortality¹⁶.

Potential dairy matrix mechanisms

The explanation for the finding that dairy foods, even those containing fat and saturated fat such as cheese, have a neutral or even a beneficial effect on CVD is likely to lie in the complex composition of the dairy matrix²³.

Although some dairy foods contain saturated fatty acids, they are also rich in nutrients and bioactive components such as **calcium, potassium, phosphorus and bioactive peptides** that may modify CVD risk through, for example, positive effects on blood pressure, weight and diabetes. In addition, dairy constituents such as bioactive peptides may have direct effects on cardiovascular parameters including blood clotting, arterial stiffness endothelial function and inflammation²⁴. This may help counter any negative effects of saturated fat in dairy on blood lipids and CVD risk. However, it is also increasingly recognised that individual **saturated fatty acids** have different effects on blood lipids; several of those in milk fat do not have an adverse effect on LDL ('bad') cholesterol or other markers of CVD risk including HDL ('good') cholesterol and the ratio of total to HDL cholesterol²⁵. Moreover, components of the dairy matrix have been shown to modify blood lipid levels²³.

There is evidence that **calcium** in dairy foods, through its effects on binding fat and decreasing its absorption in the gut, may reduce the potential rise in LDL cholesterol following saturated fat consumption²⁶⁻²⁸. For example, cheese does not increase LDL cholesterol compared with butter of equal fat content²⁷. Similarly, compared with a low-calcium control diet, milk- and cheese-based diets lessened saturated fatty acid-induced increases in total and LDL cholesterol²⁹ It may be important for this beneficial effect that fat and calcium are embedded in the same food

matrix, as is the case for milk and cheese²³. **Phosphorus** in the dairy matrix may also interact with calcium to influence blood lipids; calcium phosphate binds bile acids and fatty acids, and increases their excretion³⁰. It has been suggested too that the membrane which encloses milk fat (the **milk fat globule membrane**; MFGM) and which is rich in bioactive phospholipids and proteins may have a beneficial role in modulating blood lipids³¹. Fermented dairy foods may also modify blood lipids through favouring gut bacteria which produce short-chain fatty acids (SCFA), and which in turn have a positive effect on lipids³².

In addition, the total fatty acid profile of a food, not just its saturated fatty acid content, is important.

Milk fat includes a number of fatty acids which may have beneficial effects on CVD risk factors such as blood lipids and markers of inflammation. These include conjugated linoleic acid (cis-9, trans-11 CLA) and trans palmitoleic acid (trans-C16:1)³³⁻³⁵. A recent study reported that higher levels of a biomarker of dairy fat in the diet were associated with a lower incidence of CVD and CVD risk factors³⁶.

It is evident that in terms of the effects of milk and dairy foods on cardiovascular health, the whole food, and the dietary pattern, rather than an individual component such as saturated fat should be taken into account. In this respect, the weight of epidemiological evidence suggests no adverse effects of regularly consuming milk and dairy foods on cardiovascular health and rather, in some studies, a cardio-protective effect has been observed.

Cancer

Based on a comprehensive review of the scientific literature, the World Cancer Research Fund report on diet and cancer concludes there is strong evidence that dairy foods are likely to protect against colorectal cancer. There is also limited evidence that consuming dairy foods might decrease the risk of premenopausal breast cancer. Although a protective effect of milk and milk products on bladder cancer and postmenopausal breast cancer has been indicated, the evidence is inconclusive. The evidence suggesting an increased risk of prostate cancer with higher consumption of dairy products is limited.

Several plausible mechanisms exist whereby constituents of the dairy matrix, either individually or in combination, may be involved in protective effects including calcium, lactoferrin, conjugated linoleic acid (CLA), sphingolipids, vitamin K2, probiotics and vitamin D (in fortified dairy).

Both protective and negative associations with dairy foods and cancer have been suggested by epidemiological studies. The relationship between diet and cancer risk is complex and pooled analysis of data is necessary before a reliable picture emerges. Even then, most common cancers are multifactorial, generally resulting from several causative factors and the impact of individual foods or food groups, either positive or negative, may be relatively small. The World Cancer Research Fund (WCRF) together with the American Institute for Cancer Research (AICR) systematically analyses the worldwide scientific literature on the relationship between food, nutrition, physical activity and cancer. The reports produced include an assessment of the strength of the evidence and the mechanisms which may be involved.

Colorectal Cancer

There is strong evidence that consuming dairy products decreases the risk of colorectal cancer

(cancer of the colon and rectum, sometimes referred to as bowel cancer). Meta-analyses consistently report a reduced risk with higher intakes of milk and dairy foods. The latest WCRF/AICR Report (2017) concludes that consumption of dairy foods probably protects against colorectal cancer, and rates the level of evidence as 'strong'¹. The report estimates a 13% decreased risk of colorectal cancer per 400g of dairy consumed a day. A significant dose-response relationship was also noted with milk: a 6% decreased risk of colorectal cancer per 200g of milk consumed a day.

The protective association is observed in European populations, as well as North American. For example, results from the European Prospective Investigation into Cancer and Nutrition (EPIC) which included data from over 477,000 men and women from ten European countries showed that higher intakes of milk, cheese, yogurt and total dairy products (and dietary calcium from dairy sources) were all associated with reduced colorectal cancer risk².

Bladder Cancer

It has been suggested that milk intake may have a protective effect on bladder cancer. However, according to the 2015 WCRF report, evidence for an association is inconsistent and no conclusions could be drawn³. A meta-analysis in 2011 showed a decreased risk of bladder cancer with a high intake of milk, while another in the same year found no association between dairy intake and bladder cancer risk^{4, 5}.

Breast cancer

The most recent WCRF/AICR report (2017) concludes that there is limited, but generally consistent, evidence that consumption of dairy products may decrease the risk of premenopausal breast cancer⁶. A dose response meta-analysis conducted as part of the report estimated a 5% reduced risk of breast cancer per 400g of dairy foods consumed a day. Evidence for an association between dairy intake and risk of postmenopausal breast cancer was inconclusive.

Prostate cancer

The 2014 WCRF report concluded that the evidence of an association between dairy intake and increased risk of prostate cancer is limited^{7,8}. The evidence for total dairy product intake showed an increased risk per 400g a day, but the relationship was unclear and not significant when stratified by prostate cancer type. For diets high in calcium, WCRF also concludes that the evidence suggesting an increased risk of prostate cancer is limited⁷. On this basis, dairy is not part of the WCRF dietary recommendations concerning prostate cancer risk⁷.

Other cancers

In relation to other cancer types including ovarian, endometrial, lung and pancreatic cancers, WCRF reports and recent meta-analyses suggest no association with dairy intake⁹. In Europe and the United States, there is some suggestion that total dairy product intake may reduce the risk of gastric cancer¹⁰.

Potential dairy matrix mechanisms

Dairy foods contain several components which may have anti-cancer properties. The effect of milk, or other dairy foods, in reducing colorectal cancer risk is likely to be mediated, at least in part, by their **calcium** content. Plausible mechanisms include calcium's ability to bind secondary bile acids and free fatty acids in the gut lumen which can otherwise have a toxic effect on the cells of the colon, and to reduce abnormal proliferation of colonic epithelial cells¹¹. Similarly, animal studies have shown that a high calcium intake also inhibits hyper-proliferation of mammary glands and has inhibitory effects on breast cancer cells¹². Indeed, a 2016 meta-analysis reports a dose-response relationship between calcium intake and reduced breast cancer risk¹³.

There is evidence too that other components of the dairy matrix may be involved along with calcium in protective effects. For example, the relation between dietary calcium and a lower risk of colorectal cancer in the EPIC study was evident for dairy sources of calcium only¹⁴. Similarly, in a prospective cohort of Swedish men, control for total calcium intake in the analysis lessened but did not eliminate the beneficial effect of milk¹⁵.

Other constituents of milk and milk products which may have anti-carcinogenic properties include **vitamin D** (in fortified dairy), **lactoferrin** (a protein in cows' milk), **menaquinones** (a class of **vitamin K** compounds of which cheese is an important dietary source) and **probiotic** bacteria in fermented products such as

yogurt¹⁶⁻²⁰. In addition, the fatty acids **conjugated linoleic acid** (CLA) and **butyric acid**, and components of the membrane which encloses the fat droplets in milk (milk fat globule membrane; MFGM) such as **sphingolipids** and particularly sphingomyelin, are thought to have anti-cancer effects^{16, 20-22}. With respect to CLA, data from the Swedish Mammography Cohort found intakes of CLA could partly explain the relation between high-fat dairy food and lower colorectal cancer incidence observed in this study²³. However, in relation to other cancers, including breast cancer, although promising from in vitro and animal work, the evidence for CLA from human studies is limited²⁴. The mechanisms by which dairy constituents may reduce cancer risk are not mutually exclusive and there may be interaction within the dairy matrix to produce the beneficial effects.

Research on the role of specific dairy products and dairy constituents, and possible interactions between them in the matrix will help clarify our understanding of the relationship between dairy and cancer risk. Nevertheless, from the available data there is good evidence that milk reduces the risk of colon cancer. In addition, milk and milk products may also be associated with a reduced risk of premenopausal breast cancer and possibly bladder cancer. Consumption of milk and dairy foods as part of a healthy dietary pattern fits with guidance on cancer prevention²⁵.

Muscle mass maintenance in older people

There is evidence to suggest a potential role for milk and dairy foods in helping to maintain muscle mass and function in older people. A number of studies point to the benefits of milk protein for increasing muscle protein synthesis in the elderly, and that supplementation, in combination with physical activity, can improve muscle mass and function. There is some evidence too, that older people with higher intakes of milk, cheese and yogurt have greater muscle mass and better functional capacity. In addition to high-quality protein for muscle health, the rich mix of other nutrients in milk and dairy foods make them a valuable part of the diets of older people.



Sarcopenia

Ageing is accompanied by a progressive loss of skeletal muscle mass and strength – sarcopenia - which leads to the loss of functional capacity and a greater risk of developing metabolic disease such as diabetes¹. Although some degree of sarcopenia is inevitable, the extent to which it can be minimised, and possibly reversed, has important implications as loss of physical function predicts loss of independence, falls, and even mortality. With an ageing population in Europe, strategies for prevention are increasingly important.

Muscle protein synthesis

Both food intake and resistance exercise stimulate protein synthesis in muscles. Recent studies suggest that older people's muscles are less responsive to the stimulating effects of protein than their younger counterparts². Consequently, research has focused on whether higher intakes of protein can overcome this 'anabolic resistance' and can enhance the effects of exercise. Although there is not yet consensus, it has been argued that to help older people maintain and regain lean body mass and function, higher protein intakes than currently recommended (0.8 g/kg/day) are required: in the range of at least 1.0 to 1.2 g/kg/day and up to 1.5g/kg/day³⁻⁶. Roughly equal distribution of protein intake at meals across the day is suggested to be the most effective way to achieve this which, assuming three meals a day, equates to 0.4 to 0.5 g/kg per meal⁷. Currently, protein intakes at breakfast time are often low, and increasing protein intake before bed may also represent an opportunity for overnight muscle protein synthesis^{8,9}.

The quality of protein intake is also important. Protein that has a high concentration of essential amino acids, most importantly leucine, has been shown to best stimulate muscle protein synthesis⁹. This points to milk protein, particularly whey, and a number of studies have confirmed beneficial effects on muscle protein synthesis^{10,11}. It is likely

that the effects of whey on muscle gain in older people extend beyond its leucine or essential amino acid content since comparable amino acid 'mixes' do not have the same effect¹¹. Factors such as rate of absorption, influenced by the dairy food matrix, may be important¹²⁻¹⁴. In this respect, more research is needed on the effects of milk per se and of other whole dairy foods on muscle protein synthesis in older people.

Dairy protein supplementation

A meta-analysis in 2012 of longer term studies examining the effects of diet and exercise in older people found that protein supplementation increased muscle mass and strength gains during resistance exercise programmes: 38% more fat free mass and a 33% increase in strength¹⁵. All six studies in the meta-analysis used a dairy-based protein; five exclusively dairy (milk, whey or casein) and the sixth a combination of egg, meat and dairy. A subsequent six-month clinical trial from the Netherlands also found that a milk protein drink combined with a resistance exercise programme significantly increased skeletal muscle mass in frail elderly adults¹⁶. Another long-term study by the same research group found that although increasing milk protein without exercise did not increase muscle mass, it enhanced physical performance in the frail elderly subjects including improvements in balance, walking speed and ability to 'get up and go' from a chair¹⁷. The milk drink used in these two studies provided around 30g of protein a day, equivalent to 3 to 4 servings of dairy. In another dairy intervention, adding 210g of ricotta cheese a day to the usual diets of older people (without sarcopenia) for 12 weeks improved skeletal muscle mass and balance-test scores¹⁸.

Observational studies

There is some evidence that older people with higher intakes of dairy have greater muscle mass and better muscle function. In a prospective cohort study of older Spanish adults (60 years plus), higher consumption of low-


fat milk and yogurt was associated with lower risk of frailty and, specifically, of slow walking speed and weight loss¹⁹. Similarly, a recent study of almost 4,000 older people in Ireland (over 60 years), found higher daily yogurt intake was associated with better physical function scores²⁰. A cross-sectional study in 70 to 85-year-old Australian women also reports that those with the highest milk, cheese and yogurt consumption (2.2 or more servings a day) had significantly

greater lean body and skeletal muscle mass, greater hand-grip strength and better 'up and go' performance than women who consumed the least (less than 1.5 servings)²⁰. The authors highlight the bioactive compounds present in dairy such as high-quality proteins and interactions with other components of the dairy matrix such as calcium which may be responsible for the beneficial effects.

Ensuring adequate protein intake, including with milk protein, alongside physical activity, appears a promising approach to maintaining and improving muscle mass and functional performance in older people. Given the consequences of sarcopenia for health and quality of life, and with an ageing population, such strategies are increasingly important. In addition to protein, milk and dairy foods offer older people other valuable nutrients in a palatable, convenient and affordable way.

Recovery after exercise

Although this is a relatively new area of dairy research, milk shows promise in the sports nutrition arena. The nutritional composition of milk makes it particularly well suited to support recovery after exercise. There is evidence that milk can be an effective post-exercise rehydration drink due to its fluid and electrolyte content. The protein in milk also helps promote muscle protein synthesis after exercise, and milk has been shown to reduce exercise-induced muscle damage and soreness. Practically, milk is convenient, inexpensive and accessible.



The nutritional composition of the milk matrix, including its protein, carbohydrate and electrolyte content suggests that it would be useful as a sports drink. A number of studies have confirmed a potential role for milk in sports and exercise nutrition, particularly in relation to recovery after exercise¹.

Rehydration

During exercise, fluid is lost from the body as sweat and needs to be replaced. The main factors that influence the process of post-exercise rehydration are the volume and composition of the fluid consumed, particularly the electrolyte concentration. The **sodium** and **potassium** content of milk make it a good candidate for effective post-exercise rehydration, and several studies have now

shown that low-fat milk can restore and maintain hydration status equally as well as, or better than, a commercially-available sports drink²⁻⁵. The rehydration potential has been demonstrated in children and teens as well as adults⁶. In addition to the electrolyte content, there is some evidence that the **protein** in milk may also enhance rehydration, possibly through slowed gastric emptying⁷. Again, this has been shown in children as well as adults⁸. A recent trial to assess the potential of 12 different drinks to affect hydration status reports that milk (skimmed and whole milk) and an oral rehydration solution were the most effective at maintaining fluid balance⁹.

Muscle recovery and repair

Following exercise, protein is important for recovery and repair.

Resistance exercise stimulates muscle protein synthesis but a net gain in muscle mass is only possible if adequate protein or essential amino acids are also consumed. Milk is rich in high quality protein (80% casein and 20% whey) and contains all the essential amino acids; it is a good source of branched chain amino acids including leucine which are integral to muscle metabolism and produces a sustained increase in blood amino acids. Studies support a beneficial effect of milk and of dairy proteins in recovery from resistance exercise. Milk can stimulate protein synthesis and support muscle development following bouts of resistance exercise, in men and women, and in the short and longer term¹⁰⁻¹³ and may have advantages for muscle metabolism over other protein

sources such as soy^{10,12}. Studies of individual milk proteins, particularly whey, also support beneficial effects on skeletal muscle amino acid uptake, protein synthesis and muscle mass¹⁴. In relation to muscle repair after exercise, studies have shown that drinking milk immediately following resistance exercise (500ml) can help to alleviate muscle soreness and drops in muscle performance¹⁵⁻¹⁸.

Glycogen re-synthesis

Carbohydrate is key to supporting glycogen re-synthesis after exercise.

Milk contains carbohydrate as the naturally present sugar lactose (glucose and galactose) and so can contribute to glycogen re-synthesis. Studies in this area have largely focused on flavoured milk, particularly chocolate milk, which has greater amounts of carbohydrate and has been shown to be effective for post-exercise muscle glycogen recovery^{1,19}.

Calcium balance

A further area where milk and dairy foods may be of benefit in sports nutrition is in relation to calcium balance.

It has been hypothesised that calcium loss through prolonged or excessive sweating may have a detrimental effect on bone²⁰. Although more research is needed, a recent study in competitive female cyclists found that a dairy-based meal before exercise was able to counteract the loss of calcium in sweat and reduce bone breakdown²¹.

It is well established that milk can be effective for rehydration, and for muscle recovery and repair after exercise and sport. Preliminary data suggests it may also be of value pre-exercise as a calcium provider. Practically, milk is convenient, inexpensive and accessible. While chocolate milk and whey proteins also have a role in sports nutrition, the potential of other dairy products such as yogurt and cheese in this context has yet to be fully elucidated. The rich nutrient content of these dairy foods, however, mean they make a valuable contribution to a healthy, balanced diet for sports people.



Conclusion

Milk and dairy foods are naturally rich sources of a wide range of nutrients and make a significant contribution to nutrient intakes and diet quality in Europe. The milk and dairy food group forms an important part of food-based dietary guidelines across Europe, and dairy is, with good reason, recognised as one of the components of a healthy dietary pattern. The specific recommendations vary between countries, but on average 2 to 3 servings of milk and dairy foods a day are recommended for adults; often more for children and adolescents (3 to 4 servings), and in some cases more for pregnant women and older people too.

As this review of the research indicates, consumption of milk and dairy foods is linked to a number of potential health benefits including for bone, blood pressure, weight control, type 2 diabetes, cardiovascular disease and colorectal cancer. They also have a role in sports nutrition and in the helping to maintain muscle mass and function in older people. It is increasingly recognised that the unique combination of nutrients and bioactive factors, and how they interact with each other in the dairy matrix, combine to produce these positive effects on health.

Milk and dairy foods are versatile and offer a wide range of nutritional benefits in an enjoyable, convenient and affordable way.





References

Dairy food consumption data sources

Austria:

Austrian Nutrition Report 2008

Figures refer to adults 18 – 65 years.

Elmadfa I, Freisling H, Nowak V, Hofstätter D et al. Vienna, March 2009

Belgium:

Food Consumption Survey, Belgium 2014-2015

Figures refer to population 3-64 years.

Total dairy = cheese, milk, yogurt, milk deserts (e.g. pudding), milk drinks, curd (~ fresh white cheese), soy drinks, soy deserts De Ridder K, Bel S, Brocatus L, Lebacq T, Moyersoen I, Ost C & Teppers E. De consumptie van voedings-middelen en de inname van voedingsstoffen. In: Bel S, Tafforeau J (ed.). Voedselconsumptiepeiling 2014-2015. Rapport 4. WIV-ISP, Brussels, 2016

Denmark:

Dietary habits in Denmark 2011-2013

Figures refer to the total population, rather than adults only.

Main results: DTU Food – National Food Institute. Copenhagen 2015

France:

INCA 3: Third National Nutrition Survey 2014-2015

Figures refer to percentage contribution of 'milk, yogurt, cottage cheese, cheese and milk desserts' food group for French adults aged 18-79 years.

ANSES (French Agency for Food, Environmental and Occupational Health & Safety)

Detailed results of INCA3 survey:

<https://www.anses.fr/fr/system/files/NUT2014SA0234Ra.pdf> (report in French, June 2017)

Ireland:

National Adult Nutrition Survey 2008-2010

Figures refer to percentage contribution of all milk, cheese, yogurt, cream and butter for Irish adults aged 18-90 years.

Feeney E et al. An overview of the contribution of dairy and cheese intakes to nutrient intakes in the Irish diet: Results from the National Adult Nutrition Survey. Br J Nutr. 2016; 115: 709-717.

Netherlands:

Dutch National Food Consumption Survey 2007-2010

Diets of children and adults aged 7 to 69 years, National Institute for Public Health and the Environment 2011. Report number 350050006/2011.

Report Dairy Consumption in the Netherlands. Results of the Dutch National Food Consumption Survey 2007-2010. Report written by Dr Diewertje Sluik and Prof Dr Edith Feskens, July 2013 (this Report is in Dutch).

Norway:

National diet surveys of adults from 2010 (Norkost3).

Figures refer to percentage contribution of milk, yogurt, cheese, butter and cream products for population aged 18-70 years to nutrient intakes. Survey conducted by Department of Nutrition, University of Oslo, the Directorate of Health, and the Norwegian Food Safety Authority.

Risiko for jodmangel i Norge. Identifisering av et akutt behov for tiltak. Nasjonalt råd for ernæring, Helsedirektoratet, IS-0591 (this report is in Norwegian, June 2016).

UK:

National Diet and Nutrition Survey 2012/13 – 2013/14

Figures refer to percentage contribution of the 'milk, yogurt & cheese' food group for UK adults aged 19-64 years.

National Diet and Nutrition Survey. Results from Years 5 and 6 (combined) of the Rolling Programme (2012/13 – 2013/14). A survey carried out on behalf of Public Health England and the Food Standards Agency, 2016

Nutrient richness

1. European Commission. EU Register of nutrition and health claims made on foods EU Register on nutrition and health claims. <http://ec.europa.eu/nuhclaims/> (accessed 11/10/13).

2. Vissers PA et al. Contribution of dairy products to micronutrient intake in The Netherlands. J Am Coll Nutr. 2011; 30(5 Suppl 1): 415S-421S.

3. Coudray B. Contribution of dairy products to micronutrient intake in France. J Am Coll Nutr. 2011; 30(5 Suppl 1): 410S-414S.

4. Fulgoni VL 3rd et al. Nutrients from dairy foods are difficult to replace in diets of Americans: food pattern modeling and an analyses of the National Health and Nutrition Examination Survey 2003-2006. Nutr Res. 2011; 31: 759-765.

5. Mensink GB et al. Mapping low intake of micronutrients across Europe. *Br J Nutr.* 2013; 110: 755–773.
 6. Gregory J et al. *The National Diet and Nutrition Survey: Young People aged 4 to 18 years, Volume 1: Report of the Diet and Nutrition Survey.* London: TSO, 2000.
 7. Nordic Council of Ministers. *Nordic Nutrition Recommendations 2012 - Part 1. Summary, Principles and Use.* 5th edition. Copenhagen, Denmark: Nordic Council of Ministers; 2013. <http://www.norden.org/en/publications/publikationer/nord-2013-009> (accessed 20/10/2013).
 8. Drewnowski A & Fulgoni V 3rd. Nutrient profiling of foods: creating a nutrient-rich food index. *Nutr Rev.* 2008; 66: 23-39.
 9. Drewnowski A & Darmon N. The economics of obesity: dietary energy density and energy cost. *Am J Clin Nutr.* 2005; 82: 265S-273S.
 10. Maillot M et al. Nutrient profiling can help identify foods of good nutritional quality for their price: a validation study with linear programming. *J Nutr.* 2008; 138: 1107-1113.
 11. Drewnowski A et al. Low-energy-density diets are associated with higher diet quality and higher diet costs in French adults. *J Am Diet Assoc.* 2007; 107: 1028-1032.
 12. Darmon N & Drewnowski A. The contribution of food prices and diet cost to socioeconomic disparities in diet quality: a systematic review and analysis. *Nutr Rev.* 2015; 7: 643-660.
 13. De Ridder K, Bel S, Brocatus L, Lebacqz T, Moyersoelen I, Ost C & Teppers E. De consumptie van voedingsmiddelen en de inname van voedingsstoffen. In: Bel S, Tafforeau J (ed.). *Voedselconsumptiepeiling 2014-2015. Rapport 4.* WIV-ISP, Brussel, 2016
 14. Irish Universities Nutrition Alliance. *Report on the Contribution of Dairy Foods to the Nutritional Quality of the Irish Adult Diet* Commissioned by the National Dairy Council through Funds Awarded from the Dairy Research Trust Co-operative Society Ltd, 2012. Analysis based on the National Adult Nutrition Survey (NANS).
 15. Laird E et al. Dairy intakes in older Irish adults and effects on vitamin micronutrient status: data from the TUDA study. *J Nutr Health Aging.* 2016. doi:10.1007/s12603-016-0845-4
 16. CCAF (French eating behaviours and consumption) Surveys 2013 and 2016, CREDOC
 17. Health Council of the Netherlands. *Dutch dietary guidelines 2015.* The Hague: Health Council of the Netherlands, 2015; publication no. 2015/24E.
- Dairy matrix effects**
1. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.
 2. Chen GC et al. Cheese consumption and risk of cardiovascular disease: a meta-analysis of prospective studies. *Eur J Nutr.* 2016 Aug 12; DOI: 10.1007/s00394-016-1292-z.
 3. Muehlhoff et al. *Milk and dairy products in human nutrition.* Rome: Food and Agriculture Organisation of the United Nations; 2013. <http://www.fao.org/docrep/018/i3396e/i3396e.pdf> (accessed 01/12/13).
 4. St-Onge MP et al. Consumption of fermented and nonfermented dairy products: effects on cholesterol concentrations and metabolism. *Am J Clin Nutr.* 2000; 71: 674–681.
 5. Parodi PW. Milk fat in human nutrition. *Aust J Dairy Technol.* 2004; 59: 3-59.
 6. Rosqvist F et al. Potential role of milk fat globule membrane in modulating plasma lipoproteins, gene expression, and cholesterol metabolism in humans: a randomized study. *Am J Clin Nutr.* 2015; 102: 20-30.
 7. Tsuchiya A et al. Higher satiety ratings following yogurt consumption relative to fruit drink or dairy fruit drink. *J Amer Diet Assoc.* 2006; 106: 550-557.
 8. Churchward-Venne TA et al. Ingestion of casein in a milk matrix modulates dietary protein digestion and absorption kinetics but does not modulate postprandial muscle protein synthesis in older men. *J Nutr.* 2015; 145: 1438-1445.

Bone health

1. Heaney RP. Dairy and bone health. *J Am Coll Nutr.* 2009; 28: 82S-90S.
2. Dietary Guidelines Advisory Committee. Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2010. Washington, DC: US Department of Agriculture, Agricultural Research Service; 2010. <http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2010/DGAC/Report/2010DGACReport-camera-ready-Jan11-11.pdf> (accessed 11/12/13).
3. Rozenberg S et al. Effects of dairy product consumption on health: benefits and beliefs - a commentary from the Belgian Bone Club and the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases. *Calcif Tissue Int.* 2016; 98: 1–17.
4. Huncharek M et al. Impact of dairy products and dietary calcium on bone-mineral content in children: results of a meta-analysis. *Bone.* 2008; 43: 312-321.
5. van den Hooven EH et al. Identification of a dietary pattern prospectively associated with bone mass in Australian young adults. *Am J Clin Nutr.* 2015; 102: 1035-1043.
6. Bonjour JP et al. Gain in bone mineral mass in prepubertal girls 3½ years after discontinuation of calcium supplementation: a follow-up study. *Lancet.* 2001; 358: 1208-1212.
7. Cheng S et al. Effects of calcium, dairy product, and vitamin D supplementation on bone mass accrual and body composition in 10–12-y-old girls: a 2-y randomized trial. *Am J Clin Nutr.* 2005; 82: 1115–1126.
8. Cadogan J et al. Milk intake and bone mineral acquisition in adolescent girls: randomized controlled intervention trial. *BMJ.* 1997; 315: 1255-1260.
9. Weaver CM. Milk Consumption and Bone Health. *JAMA Pediatr.* 2014; 168: 12-13.
10. Black RE et al. Children who avoid drinking cow milk have low dietary calcium intakes and poor bone health. *Am J Clin Nutr.* 2002; 76: 675-680.
11. Goulding A et al. Children who avoid drinking cow's milk are at increased risk for prepubertal bone fractures. *J Am Diet Assoc.* 2004; 104: 250-253.
12. Weaver CM et al. The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. *Osteoporos Int.* 2016; 27: 1281-1386.
13. Heaney RP. Calcium, dairy products and osteoporosis. *J Am Coll Nutr.* 2000; 19: 83S-99S.
14. Manios Y et al. Changes in biochemical indexes of bone metabolism and bone mineral density after a 12-mo dietary intervention program: the Postmenopausal Health Study. *Am J Clin Nutr.* 2007; 86: 781-789.
15. Moschonis G et al. The effects of a 30-month dietary intervention on bone mineral density: the Postmenopausal Health Study. *Br J Nutr.* 2010; 104:100-107.
16. Bischoff-Ferrari HA et al. Milk intake and risk of hip fracture in men and women: a meta-analysis of prospective cohort studies. *J Bone Miner Res.* 2011; 26: 833-839.
17. Michaëlsson K et al. Milk intake and risk of mortality and fractures in women and men: cohort studies. *BMJ.* 2014; 349:g6015
18. Honkanen R et al. Lactose intolerance associated with fractures of weight-bearing bones in Finnish women aged 38-57 years. *Bone.* 1997; 21: 473-477.
19. Tang BM et al. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet.* 2007; 370: 657-666.
20. Guéguen L & Pointillart A. The bioavailability of dietary calcium. *J Am Coll Nutr.* 2000; 19: 119-136.
21. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.
22. Kerstetter JE. Do dairy products improve bone density in adolescent girls? *Nutr Rev.* 1995; 53: 328-332.

23. Surdykowski AK et al. Optimizing bone health in older adults: the importance of dietary protein. *Aging Health*. 2010; 6: 345-357.

24. Bonjour JP. Protein intake and bone health. *Int J Vitam Nutr Res*. 2011; 81: 134-142.

25. Bonjour JP et al. Dairy in adulthood: from foods to nutrient interactions on bone and skeletal muscle health. *J Am Coll Nutr*. 2013; 32: 251-263.

26. Rizzoli R et al. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: A consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). 2014; *Maturitas* 17 July 2014;doi: 10.1016/j.maturitas.2014.1007.1005. doi:10.1016/j.maturitas.2014.07.005.

27. Kerstetter, JE et al. Dietary protein and skeletal health: a review of recent human research. *Curr Opin Lipidol*. 2011; 22:16-20.

28. Fenton TR et al. Causal assessment of dietary acid load and bone disease: a systematic review & meta-analysis applying Hill's epidemiologic criteria for causality. *Nutr J*. 2011; 10: 41-64.

29. Mangano KM et al. Dietary protein is beneficial to bone health under conditions of adequate calcium intake: an update on clinical research. *Curr Opin Clin Nutr Metab Care*. 2014; 17: 69-74.

30. Langsetmo L et al. The association between protein intake by source and osteoporotic fracture in older men: a prospective cohort study. *J Bone Miner Res*. 2016 Dec 12. doi: 10.1002/jbmr.3058. [Epub ahead of print]

31. Durosier-Izart C et al. Peripheral skeleton bone strength is positively correlated with total and dairy protein intakes in healthy postmenopausal women. *Am J Clin Nutr*. 2017; doi: 10.3945/ajcn.116.134676

32. McCabe L et al. Prebiotic and probiotic regulation of bone health: role of the intestine and its microbiome. *Curr Osteoporos Rep*. 2015; 13: 363–371.

33. Bonjour JP. Calcium and phosphate: a duet of ions playing for bone health. *J Am Coll Nutr*. 2011; 30: 438S-448S.

Blood pressure

1. Livingstone KM et al. Does dairy food intake predict arterial stiffness and blood pressure in men? Evidence from the Caerphilly Prospective Study. *Hypertension*. 2013; 61: 42-47.

2. Engberink MF et al. Inverse association between dairy intake and HTN: The Rotterdam Study. *Am J Clin Nutr*. 2009; 89: 1877-1883.

3. Engberink MF et al. Dairy intake, blood pressure, and incident HTN in a general Dutch population. *J Nutr*. 2009; 139: 582-587.

4. Fumeron F et al. Dairy products and the metabolic syndrome in a prospective study, DESIR. *J Am Coll Nutr*. 2011; 30(5 Suppl 1): 454S-463S.

5. Ruidavets JB et al. Independent contribution of dairy products and calcium intake to blood pressure variations at a population level. *J Hypertens*. 2006; 24: 671-681.

6. Feeney EL et al. Patterns of dairy food intake, body composition and markers of metabolic health in Ireland: results from the National Adult Nutrition Survey. *Nutr Diabetes*. 2017; 7: e243; doi:10.1038/nutd.2016.54.

7. Ralston RA et al. A systematic review and meta-analysis of elevated blood pressure and consumption of dairy foods. *J Hum Hypertens*. 2012; 26: 3-13.

8. Sodamah-Muthu SS et al. Dairy consumption and incidence of hypertension: a dose response meta-analysis of prospective cohort studies. *Hypertension*. 2012; 60: 1131-1137.

9. Dietary Guidelines Advisory Committee. Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2010. Washington, DC: US Department of Agriculture, Agricultural Research Service; 2010. <http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2010/DGAC/Report/2010DGACReport-camera-ready-Jan11-11.pdf> (accessed 11/12/13).

10. Drouin-Chartier J-P et al. Systematic review of the association between dairy product consumption and risk of cardiovascular-related clinical outcomes. *Adv Nutr*. 2016; 7: 1026–1040.

11. Appel LJ et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med.* 1997; 336: 1117-1124.
12. Sacks FM et al. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med.* 2001; 344: 3-10.
13. Siervo M et al. Effects of the Dietary Approach to Stop Hypertension (DASH) diet on cardiovascular risk factors: a systematic review and meta-analysis. *Br J Nutr.* 2015; 113: 1-15.
14. US Department of Health and Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute. Your guide to lowering your blood pressure with DASH (NIH Publication No. 06-4082). Bethesda, Maryland: NIH; 2006. http://www.nhlbi.nih.gov/health/public/heart/hbp/dash/new_dash.pdf (accessed 6/12/13).
15. Appel LJ et al. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA.* 2003; 289: 2083-2093.
16. Harnden KE et al. Dietary Approaches to Stop Hypertension (DASH) diet: applicability and acceptability to a UK population. *J Hum Nutr Diet.* 2010; 23: 3-10.
17. Mancia G et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* 2013; 31: 1281–1357.
18. Moore LL et al. Intake of fruits, vegetables, and dairy products in early childhood and subsequent blood pressure change. *Epidemiology.* 2005; 16: 4-11.
19. Asghari G et al. Dietary Approaches to Stop Hypertension (DASH) dietary pattern is associated with reduced incidence of metabolic syndrome in children and adolescents. *J Pediatr.* 2016; 174: 178-184.
20. Chiu S et al. Comparison of the DASH (Dietary Approaches to Stop Hypertension) diet and a higher-fat DASH diet on blood pressure and lipids and lipoproteins: a randomized controlled trial. *Am J Clin Nutr.* 2016; 103: 341-347.
21. Drouin-Chartier JP et al. Impact of dairy consumption on essential hypertension: a clinical study. *Nutr J.* 2014; 13: 83.
22. Machin DR et al. Hypotensive effects of solitary addition of conventional nonfat dairy products to the routine diet: a randomized controlled trial. *Am J Clin Nutr.* 2014; 100: 80-87.
23. Crippa G et al. Os 04-04 Antihypertensive effect of milk-derivative tripeptides. Randomized, double-blind, placebo-controlled study on the effects of Grana Padano cheese dop in hypertensive patients. *J Hypertens.* 2016; 34 Suppl 1: e55-56.
24. Ricci I et al. Milk protein peptides with angiotensin I-converting enzyme inhibitory (ACEI) activity. *Crit Rev Food Sci Nutr.* 2010; 50: 390-402.
25. Fekete AA et al. Casein-derived lactotripeptides reduce systolic and diastolic blood pressure in a meta-analysis of randomised clinical trials. *Nutrients.* 2015; 7: 659-681.
26. Neilsen R et al. Short communication: Is consumption of a cheese rich in angiotensin-converting enzyme-inhibiting peptides, such as the Norwegian cheese Gamalost, associated with reduced blood pressure? *J Dairy Sci.* 2014; 97: 2662–2668.
27. Fekete AA et al. Whey protein lowers blood pressure and improves endothelial function and lipid biomarkers in adults with prehypertension and mild hypertension: results from the chronic Whey2Go randomized controlled trial. *Am J Clin Nutr.* 2016; 104: 1534-1544.
28. Wilson CP et al. Riboflavin offers a targeted strategy for managing hypertension in patients with the MTHFR 677TT genotype: a 4-y follow-up. *Am J Clin Nutr.* 2012; 95: 766-772.
29. McNulty et al. Riboflavin lowers homocysteine in individuals homozygous for the MTHFR 677C->T polymorphism. *Circulation.* 2006; 113: 74-80.
30. Houston MC & Harper KJ. Potassium, magnesium, and calcium: their role in both the cause and treatment of hypertension. *J Clin Hypertens.* 2008; 10: 3-11.
31. McGrane MM et al. Dairy consumption, blood pressure, and risk of hypertension: An evidence-based review of recent literature. *Curr Cardiovasc Risk Rep.* 2011; 5: 287-298.

32. Alonso A et al. Dietary phosphorus, blood pressure, and incidence of hypertension in the atherosclerosis risk in communities study and the multi-ethnic study of atherosclerosis. *Hypertension*. 2010; 55: 776-784.

33. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.

Weight control

1. McCarron DA et al. Blood pressure and nutrient intake in the United States. *Science*. 1984; 224: 1392-1398.

2. Zemel MB et al. Regulation of adiposity by dietary calcium. *FASEB J*. 2000; 14: 1132-1138.

3. Heaney RP & Rafferty K. Preponderance of the evidence: an example from the issue of calcium intake and body composition. *Nutr Rev*. 2009; 67: 32-39.

4. Van Loan M. The role of dairy foods and dietary calcium in weight management. *J Am Coll Nutr*. 2009; 28: 120S-129S.

5. Astrup A et al. Dairy beverages and energy balance. *Physiol Behav*. 2010; 100: 67-75.

6. Dougkas A et al. Associations between dairy consumption and body weight a review of the evidence and underlying mechanisms. *Nutr Res Rev*. 2011; 24: 72-95.

7. Wang W et al. Association of dairy products consumption with risk of obesity in children and adults: a meta-analysis of mainly cross-sectional studies. *Ann Epidemiol*. 2016; 22 pii: S1047-2797.

8. Varenna M et al. Effects of dietary calcium intake on body weight and prevalence of osteoporosis in early postmenopausal women. *Am J Clin Nutr*. 2007; 86: 639–644.

9. Crichton GE, Alkerwi A. Whole fat dairy food intake is inversely associated with obesity prevalence: findings from the Observation of Cardiovascular Risk Factors in Luxembourg study. *Nutr Res*. 2014 ;34: 936-943.

10. Feeney EL et al. Patterns of dairy food intake, body composition and markers of metabolic health in Ireland: results from the National Adult Nutrition Survey. *Nutr Diabetes*. 2017; 7: e243; doi:10.1038/nutd.2016.54.

11. Louie JC et al. Dairy consumption and overweight and obesity: a systematic review of prospective cohort studies. *Obes Rev*. 2011; 12: e582-e592.

12. Rosell M et al. Association between dairy food consumption and weight change over 9 y in 19 352 perimenopausal women. *Am J Clin Nutr*. 2006; 84: 1481-1488.

13. Vergnaud AC et al. Dairy consumption and 6-y changes in body weight and waist circumference in middle-aged French adults. *Am J Clin Nutr*. 2008; 88: 1248-1255.

14. Schwingshackl L et al. Consumption of dairy products in relation to changes in anthropometric variables in adult populations: a systematic review and meta-analysis of cohort studies. *PLoS One*. 2016 Jun 16;11(6):e0157461

15. Kratz M et al. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr*. 2013; 52: 1-24.

16. Spence LA et al. The role of dairy products in healthy weight and body composition in children and adolescents. *Curr Nutr Food Sci*. 2011; 7: 40-49.

17. Bigornia SJ et al. Dairy intakes at age 10 years do not adversely affect risk of excess adiposity at 13 years. *J Nutr*. 2014; 144: 1081-1090.

18. Lu L et al. Long-term association between dairy consumption and risk of childhood obesity: a systematic review and meta-analysis of prospective cohort studies. *Eur J Clin Nutr*. 2016; 70: 414-423.

19. Abargouei AS et al. Effect of dairy consumption on weight and body composition in adults: a systematic review and meta-analysis of randomized controlled clinical trials. *Int J Obes*. 2012; 36: 1485-1493.

20. Stonehouse W et al. Dairy intake enhances body weight and composition changes during energy restriction in 18–50-year-old adults - a meta-analysis of randomized controlled trials. *Nutrients*. 2016; 8: 394.

21. Chen M et al. Effects of dairy intake on body weight and fat: a meta-analysis of randomized controlled trials. *Am J Clin Nutr*. 2012; 96: 735-747.

22. Booth AO et al. Effect of increasing dietary calcium through supplements and dairy food on body weight and body composition: a meta-analysis of randomized controlled trials *Br J Nutr*. 2015; 114: 1013-1025.
23. Soares MJ et al. Mechanistic roles for calcium and vitamin D in the regulation of body weight. *Obes Rev*. 2012; 13: 592-602.
24. Gonzalez JT et al. Effect of calcium intake on fat oxidation in adults: a meta-analysis of randomized, controlled trials. *Obes Rev*. 2012; 13: 848-857.
25. Christensen R et al. Effect of calcium from dairy and dietary supplements on faecal fat excretion: a meta-analysis of randomized controlled trials. *Obes Rev*. 2009; 10: 475-486.
26. Lorenzen JK et al. Effect of dairy calcium or supplementary calcium intake on postprandial fat metabolism, appetite, and subsequent energy intake. *Am J Clin Nutr*. 2007; 85: 678-687.
27. Major GC et al. Calcium plus vitamin D supplementation and fat mass loss in female very low-calcium consumers: potential link with a calcium-specific appetite control. *Br J Nutr*. 2009; 101: 659-663.
28. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1-13.
29. Faghhi S. et al. Comparison of the effects of cows' milk, fortified soy milk, and calcium supplement on weight and fat loss in premenopausal overweight and obese women. *Nutr Metab Cardiovasc Dis* 2011; 21: 499-503.
30. Larsen TM et al. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med*. 2010; 363: 2102-2113.
31. McGregor RA & Poppitt SD. Milk protein for improved metabolic health: a review of the evidence. *Nutr Metab*. 2013; 10: 46.
32. Tsuji H et al. Dietary medium chain triacylglycerols suppress accumulation of body fat in a double-blind, controlled trial in healthy men and women. *J Nutr*. 2001; 131: 2853-2859.
33. St-Onge MP & Jones PJ. Physiological effects of medium-chain triglycerides: potential agents in the prevention of obesity. *J Nutr*. 2002; 132: 329-332.
34. Byrne CS et al. The role of short chain fatty acids in appetite regulation and energy homeostasis. *Int J Obes*. 2015; 39: 1331-1338.
35. Swedish Council on Health Technology Assessment. Dietary Treatment of Obesity – A systematic review of the literature. Stockholm: Swedish Council on Technology Assessment in Health Care (SBU); 2013. <http://www.sbu.se/218> (accessed 1/12/13).

Type 2 diabetes

- Pittas AG et al. Review: the role of vitamin D and calcium in type 2 diabetes. A systematic review and meta-analysis. *J Clin Endocrinol Metab*. 2007; 92: 2017-2029.
- Elwood PC et al. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids*. 2010; 45: 925-939.
- Tong X et al. Dairy consumption and risk of type 2 diabetes mellitus: a meta-analysis of cohort studies. *Eur J Clin Nutr*. 2011; 65: 1027-1031.
- Sluijs I et al. The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. *Am J Clin Nutr*. 2012; 96: 382-390.
- O'Connor LM et al. Dietary dairy product intake and incident type 2 diabetes: a prospective study using dietary data from a 7-day food diary. *Diabetologia*. Epub ahead of print 8 February 2014. DOI: 10.1007/s00125-014-3176-1.
- Diaz-Lopez A et al. Dairy product consumption and risk of type 2 diabetes in an elderly Spanish Mediterranean population at high cardiovascular risk. *Eur J Nutr*. 2015 Feb 7. [Epub ahead of print]
- Fumeron F et al. Dairy products and the metabolic syndrome in a prospective study, DESIR. *J Am Coll Nutr*. 2011; 30(5 Suppl 1): 454S-463S

8. Struijk EA et al. Dairy product intake in relation to glucose regulation indices and risk of type 2 diabetes. *Nutr Metab Cardiovasc Dis.* 2013; 23: 822-828.
9. Soedamah-Muthu SS et al. Consumption of dairy products and associations with incident diabetes CHD and mortality in the Whitehall II study. *Br J Nutr.* 2013; 109: 718-726.
10. Bergholdt HK et al. Milk intake is not associated with low risk of diabetes or overweight obesity: a Mendelian randomization study in 97,811 Danish individuals. *Am J Clin Nutr.* 2015; 102(2): 487-496.
11. Eussen SJ et al. Consumption of dairy foods in relation to impaired glucose metabolism and type 2 diabetes mellitus: the Maastricht Study. *Br J Nutr.* 2016; 115: 1453-1461.
12. Aune D et al. Dairy products and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Am J Clin Nutr.* 2013; 98: 1066-1083.
13. Gao D et al. Dairy Products Consumption and Risk of Type 2 Diabetes: Systematic Review and Dose-Response Meta-Analysis. *PLOS ONE.* 2013; 8: e73965.
14. Health Council of the Netherlands. Dutch Dietary Guidelines 2015. The Hague: Health Council of the Netherlands, 2015; publication no. 2015/24E.
15. Gijsbers L et al. Consumption of dairy foods and diabetes incidence: a dose-response meta-analysis of observational studies. *Am J Clin Nutr* 2016; 103: 1111–1124.
16. Drouin-Chartier J-P et al. Systematic review of the association between dairy product consumption and risk of cardiovascular-related clinical outcomes. *Adv Nutr.* 2016; 7: 1026–1040.
17. Rideout TC et al. Consumption of low-fat dairy foods for 6 months improves insulin resistance without adversely affecting lipids or bodyweight in healthy adults: a randomized free-living cross-over study. *Nutr J.* 2013; 12: 56-64.
18. Dugan CE et al. Increased dairy consumption differentially improves metabolic syndrome markers in male and female adults. *Metab Syndr Relat Disord.* 2014; 12: 62-69.
19. Belin RJ & He K. Magnesium physiology and pathogenic mechanisms that contribute to the development of the metabolic syndrome. *Magnes Res.* 2007; 20:107-129.
20. Dong JY et al. Magnesium intake and risk of type 2 diabetes: meta-analysis of prospective cohort studies. *Diabetes Care.* 2011; 34(9): 2116-2122.
21. McGregor RA & Poppitt SD. Milk protein for improved metabolic health: a review of the evidence. *Nutr Metab.* 2013; 10: 46.
22. Jakubowicz D & Froy O. Biochemical and metabolic mechanisms by which dietary whey protein may combat obesity and Type 2 diabetes. *J Nutr Biochem.* 2013; 24: 1-5.
23. Ricci-Cabello et al. Possible role of milk-derived bioactive peptides in the treatment and prevention of metabolic syndrome. *Nutr Rev.* 2012; 70: 241-255.
24. Comerford KB et al. Emerging evidence for the importance of dietary protein source on glucoregulatory markers and type 2 diabetes: different effects of dairy, meat, fish, egg, and plant protein foods. *Nutrients.* 2016; 8: 446.
25. Beulens JW et al. Dietary phyloquinone and menaquinones intakes and risk of type 2 diabetes. *Diabetes Care.* 2010; 33: 1699–1705.
26. Ericson U et al. Food sources of fat may clarify the inconsistent role of dietary fat intake for incidence of type 2 diabetes. *Am J Clin Nutr.* 2015; 101: 1065-1080.
27. Mozaffarian D et al. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in US adults: a cohort study. *Ann Intern Med.* 2010; 153: 790-799.
28. Mozaffarian D et al. trans-Palmitoleic acid, other dairy fat biomarkers, and incident diabetes: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr.* 2013; 97: 854–861.
29. de Souza RJ et al. Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. *BMJ.* 2015; 11; 351:h3978.

30. Frouhi NG et al. Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study. *Lancet. Diabetes Endocrinol.* 2014; 2: 810–818.

31. Hellgren LI. Phytanic acid - an overlooked bioactive fatty acid in dairy fat? *Ann N Y Acad Sci.* 2010; 1190: 42-49.

32. Nestel PJ et al. Specific plasma lipid classes and phospholipid fatty acids indicative of dairy food consumption associate with insulin sensitivity. *Am J Clin Nutr.* 2014; 99: 46-53.

33. WHO, Regional Office for Europe, Diabetes, Data and statistics. <http://www.euro.who.int/en/health-topics/noncommunicable-diseases/diabetes/data-and-statistics> (accessed 11/12/15).

Cardiovascular disease

1. Elwood PC et al. Milk drinking, ischaemic heart disease and ischaemic stroke II. Evidence from cohort studies. *Eur J Clin Nutr.* 2004; 58: 718-724.

2. Goldbohm RA et al. Dairy consumption 10-y total and cardiovascular mortality: a prospective cohort study in the Netherlands. *Am J Clin Nutr.* 2011; 93: 615-627.

3. van Aerde MA et al. Dairy intake in relation to cardiovascular disease mortality and all-cause mortality: the Hoorn Study. *Eur J Nutr.* 2013; 52: 609-616.

4. Praagman J et al. Dairy products and the risk of stroke and coronary heart disease the Rotterdam Study. *Eur J Nutr.* 2015; 54: 981-990.

5. Dalmeijer GW et al. Dairy intake and coronary heart disease or stroke—a population-based cohort study. *Int J Cardiol.* 2013; 167: 925-929.

6. Larsson SC et al. Dairy consumption and risk of stroke in Swedish women and men. *Stroke.* 2012; 43: 1775-1780.

7. Sonestedt E et al. Dairy products and its association with incidence of cardiovascular disease: the Malmö diet and cancer cohort. *Eur J Epidemiol.* 2011; 26: 609-618.

8. Michaëlsson K et al. Milk intake and risk of mortality and fractures in women and men: cohort studies. *BMJ.* 2014; 349: g6015.

9. Hellstrand S. The statistical analysis and reality. Letter to *BMJ.* 2014; 349: g6015.

10. Bergholdt HK et al. Milk intake is not associated with low risk of diabetes or overweight-obesity: a Mendelian randomization study in 97,811 Danish individuals. *Am J Clin Nutr.* 2015; 102: 487-496.

11. Bongard V et al. Association of dietary patterns with 14-year all-cause mortality and cause-specific mortality. *Eur Heart J.* 2012; 33 (S1): 609-610.

12. Mente A et al. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med.* 2009; 169: 659-669.

13. Elwood PC et al. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids.* 2010; 45: 925-939.

14. Soedamah-Muthu SS et al. Milk and dairy consumption and incidence of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2011; 93: 158-171.

15. Larson SC et al. Milk consumption and mortality from all causes, cardiovascular disease, and cancer: a systematic review and meta-analysis. *Nutrients.* 2015; 7: 7749-7763.

16. Guo J et al. Milk and dairy consumption and risk of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Eur J Epidemiol.* 2017 Apr; 32(4): 269-287. doi: 10.1007/s10654-017-0243-1.

17. Drouin-Chartier J-P et al. Systematic review of the association between dairy product consumption and risk of cardiovascular-related clinical outcomes. *Adv Nutr.* 2016; 7: 1026–1040.

18. Qin LQ et al. Dairy consumption and risk of cardiovascular disease: an updated meta-analysis of prospective cohort studies. *Asia Pac J Clin Nutr.* 2015; 24: 90-100.

19. Hu D et al. Dairy foods and risk of stroke: a meta-analysis of prospective cohort studies. *Nutr Metab Cardiovasc Dis.* 2014; 24: 460-469.

20. de Goede J et al. Dairy consumption and risk of stroke: a systematic review and updated dose-response meta-analysis of prospective cohort studies. *J Am Heart Assoc.* 2016 May 20; 5(5). doi: 10.1161/JAHA.115.002787.
21. Alexander D et al. Dairy consumption and CVD: a systematic review and meta-analysis. *Br J Nutr.* 2016; 115: 737–750.
22. Chen GC et al. Cheese consumption and risk of cardiovascular disease: a meta-analysis of prospective studies. *Eur J Nutr.* 2016 Aug 12; DOI: 10.1007/s00394-016-1292-z.
23. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.
24. Ricci-Cabello et al. Possible role of milk-derived bioactive peptides in the treatment and prevention of metabolic syndrome. *Nutr Rev.* 2012; 70: 241-255.
25. Kris-Etherton PM & Yu S. Individual fatty acid effects on plasma lipids and lipoproteins: human studies. *Am J Clin Nutr.* 1997; 65: 1628S-1644S.
26. Lorenzen JK & Astrup A. Dairy calcium intake modifies responsiveness of fat metabolism and blood lipids to a high-fat diet. *Br J Nutr.* 2011; 105: 1823-1831.
27. Soerensen KV et al. Effect of dairy calcium from cheese and milk on fecal fat excretion, blood lipids, and appetite in young men. *Am J Clin Nutr.* 2014; 99: 984-991. Epub ahead of print 12 March 2014. DOI: 10.3945/ajcn.113.077735.
28. Lorenzen JK et al. Milk minerals modify the effect of fat intake on serum lipid profile: results from an animal and a human short-term study. *Br J Nutr.* 2014; 111: 1412-1420.
29. Hjerpsted J et al. Cheese intake in large amounts lowers LDL-cholesterol concentrations compared with butter intake of equal fat content. *Am J Clin Nutr.* 2011; 94: 1479-1484.
30. Govers MJ et al. Characterization of the adsorption of conjugated and unconjugated bile acids to insoluble, amorphous calcium phosphate. *J Lipid Res.* 1994; 35: 741–748.
31. Rosqvist F et al. Potential role of milk fat globule membrane in modulating plasma lipoproteins, gene expression, and cholesterol metabolism in humans: a randomized study. *Am J Clin Nutr.* 2015; 102: 20-30.
32. St-Onge MP et al. Consumption of fermented and nonfermented dairy products: effects on cholesterol concentrations and metabolism. *Am J Clin Nutr.* 2000; 71: 674–681.
33. Tricon S et al. Opposing effects of cis-9, trans-11 and trans-10, cis-12 conjugated linoleic acid on blood lipids in healthy humans. *Am J Clin Nutr.* 2004; 80: 614-620.
34. Mozaffarian D et al. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in US adults: a cohort study. *Ann Intern Med.* 2010; 153: 790-799.
35. Kratz et al. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr.* 2013; 52: 1-24.
36. de Oliveira Otto MC et al. Biomarkers of dairy fatty acids and risk of cardiovascular disease in the multi-ethnic study of atherosclerosis. *J Am Heart Assoc.* 2013; 2: e000092.

Cancer

- World Cancer Research Fund International/American Institute for Cancer Research. Continuous Update Project Report: Diet, Nutrition, Physical Activity and Colorectal Cancer. 2017. Available at: [wcrf.org/colorectal-cancer-2017](http://www.wcrf.org/sites/default/files/CUP%20Colorectal%20Report_2017_Digital.pdf). http://www.wcrf.org/sites/default/files/CUP%20Colorectal%20Report_2017_Digital.pdf (accessed 07/09/17).
- Aune D et al. Dairy products and colorectal cancer risk: a systematic review and meta-analysis of cohort studies. *Ann Oncol.* 2012; 23: 37-45.
- World Cancer Research Fund International/American Institute for Cancer Research. Continuous Update Project report: Diet, nutrition, physical activity and bladder cancer. 2015. <http://www.wcrf.org/sites/default/files/Bladder-Cancer-2015-Report.pdf> (accessed 10/12/2016).
- Mao QQ et al. Milk consumption and bladder cancer risk: a meta-analysis of published epidemiological studies. *Nutr Cancer.* 2011; 63: 1263-1271.
- Li F et al. Milk and dairy consumption and risk of bladder cancer: a meta-analysis. *Urology.* 2011; 78: 1298-1305.

6. World Cancer Research Fund / American Institute for Cancer Research. Continuous Update Project Report. Food, Nutrition, Physical Activity and Breast Cancer. 2017. <http://www.wcrf.org/sBreast-Cancer-2017> (accessed 5/07/2017).
7. World Cancer Research Fund International/American Institute for Cancer Research Continuous Update Project Report: Diet, Nutrition, Physical Activity, and Prostate Cancer. 2014. <http://www.wcrf.org/sites/default/files/Prostate-Cancer-2014-Report.pdf> (accessed 10/11/2016).
8. Aune D et al. Dairy products, calcium, and prostate cancer risk: a systematic review and meta-analysis of cohort studies. *Am J Clin Nutr.* 2015; 101: 87-117.
9. World Cancer Research Fund International/American Institute for Cancer Research Continuous Update Project findings and reports. <http://www.wcrf.org/int/research-we-fund/continuous-update-project-findings-reports> (accessed 13/12/2016).
10. Guo Y et al. Dairy consumption and gastric cancer risk: a meta-analysis of epidemiological studies. *Nutr Cancer.* 2015; 67: 555-568.
11. Lamprecht SA & Lipkin M. Cellular mechanisms of calcium and vitamin D in the inhibition of colorectal carcinogenesis. *Ann N Y Acad Sci.* 2001; 952: 73-87.
12. Chen P et al. Meta-analysis of vitamin D, calcium and the prevention of breast cancer. *Breast Cancer Res Treat.* 2010; 121: 469-477.
13. Hidayat K et al. Meta-analysis of vitamin D, calcium intake and breast cancer risk: meta-analysis of prospective cohort studies. *Br J Nutr.* 2016; 116: 158-166.
14. Murphy N et al. Consumption of dairy products and colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *PLoS One.* 2013; 8: e72715.
15. Larsson SC et al. Calcium and dairy food intakes are inversely associated with colorectal cancer risk in the Cohort of Swedish Men. *Am J Clin Nutr.* 2006; 83: 667-673.
16. Pufulete M. Intake of dairy products and risk of colorectal neoplasia. *Nut Res Rev.* 2008; 21: 56-67.
17. Fleet JC (2006) Dairy consumption and the prevention of colon cancer: is there more to the story than calcium? *Am J Clin Nutr.* 83; 527-528.
18. Nimptsch K et al. Dietary vitamin K intake in relation to cancer incidence and mortality: results from the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Heidelberg). *Am J Clin Nutr.* 2010; 91: 1348-1358.
19. Duarte DC et al. The effect of bovine milk lactoferrin on human breast cancer cell lines. *J Dairy Sci.* 2011; 94: 66-76.
20. Lampe JW. Dairy products and cancer. *J Am Coll Nutr.* 2011; 30: 464S-470S.
21. Parodi PW. Milk fat in human nutrition. *Aust J Dairy Technol.* 2004; 59: 3-59.
22. Kritchevsky D. Antimutagenic and some other effects of conjugated linoleic acid. *Br J Nutr.* 2000; 83: 459-465.
23. Larsson SC et al. High-fat dairy food and conjugated linoleic acid intakes in relation to colorectal cancer incidence in the Swedish Mammography Cohort. *Am J Clin Nutr.* 2005; 82: 894-900.
24. Arab A et al. The effects of conjugated linoleic acids on breast cancer: A systematic review. *Adv Biomed Res.* 2016; 5: 115.
25. World Cancer Research Fund International. Cancer Prevention & Survival. Summary of global evidence on diet, weight, physical activity & what increases or decreases your risk of cancer. May 2017 edition. <http://www.wcrf.org/sites/default/files/CUP%20Summary%20Report%20May17.pdf> (accessed 22/07/17).

Muscle mass maintenance in older people

1. Cruz-Jentoft AJ et al. Sarcopenia: European consensus on definition and diagnosis. Report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* 2010; 39: 412-423
2. Breen L & Phillips SM. Skeletal muscle protein metabolism in the elderly: Interventions to counteract the 'anabolic resistance' of ageing. *Nutr Metab.* 2011; 8: 68.

3. Bauer J et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013; 14: 542-559.
4. Wolfe RR. The role of dietary protein in optimizing muscle mass, function and health outcomes in older individuals. *Br Med J.* 2012; 108: S88-S93.
5. Paddon-Jones D & Rasmussen BB. Dietary recommendations and the prevention of sarcopenia. *Curr Opin Clin Metab Care.* 2009; 12: 86-90.
6. Witard OC et al. Growing older with health and vitality: a nexus of physical activity, exercise and nutrition. *Biogerontology.* 2016; 17: 529-46.
7. Tieland M et al. Dietary protein intake in community-dwelling, frail, and institutionalized elderly people: scope for improvement. *Eur J Nutr.* 2012; 51: 173-179.
8. Groen BB et al. Intragastric protein administration stimulates overnight muscle protein synthesis in elderly men. *Am J Physiol Endocrinol Metab.* 2012; 302: 52-60.
9. Waters DL et al. Advantages of dietary, exercise-related, and therapeutic interventions to prevent and treat sarcopenia in adult patients: an update. *Clin Interv Aging.* 2010; 5: 259-270.
10. Paddon-Jones D et al. Differential stimulation of muscle protein synthesis in elderly humans following isocaloric ingestion of amino acids or whey protein. *Exp Gerontol.* 2006; 41: 215-219.
11. Phillips SM et al. The role of milk- and soy-based protein in support of muscle protein synthesis and muscle protein accretion in young and elderly persons. *J Am Coll Nutr.* 2009; 28: 343-354.
12. Katsanos CS et al. Whey protein ingestion in elderly results in greater muscle protein accrual than its constituent essential amino acid content. *Nutr Res.* 2008; 28: 651-658.
13. Churchward-Venne TA et al. Ingestion of casein in a milk matrix modulates dietary protein digestion and absorption kinetics but does not modulate postprandial muscle protein synthesis in older men. *J Nutr.* 2015; 145: 1438-1445.
14. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.
15. Cermak NM et al. Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *Am J Clin Nutr.* 2012; 96: 1454-1464.
16. Tieland M et al. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc.* 2012; 13: 713-719.
17. Tieland M et al. Protein supplementation improves physical performance in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc.* 2012; 13: 720-726.
18. Alemán-Mateo H et al. Nutrient-rich dairy proteins improve appendicular skeletal muscle mass and physical performance, and attenuate the loss of muscle strength in older men and women subjects: a single-blind randomized clinical trial. *Clin Interv Aging.* 2014; 9: 1517-1525.
19. Lana A et al. Dairy consumption and risk of frailty in older adults: a prospective cohort study. *J Am Geriatr Soc.* 2015; 63: 1852-1860.
20. Laird E et al. Greater yogurt consumption is associated with increased bone mineral density and physical function in older adults. *Osteoporos Int.* 2017; doi.org/10.1007/s00198-017-4049-5
21. Radavelli-Bagatini S et al. Association of dairy intake with body composition and physical function in older community-dwelling women. *J Acad Nutr Diet.* 2013; 113: 1669-1674.

Recovery after exercise

1. Roy BD. Milk, the new sports drink? A review. *J Int Soc Sports Nutr.* 2008; 5:15.
2. Shirreffs SM et al. Milk as an effective post-exercise rehydration drink. *Br J Nutr.* 2007; 98: 173-180.
3. Watson P et al. A comparison of the effects of milk and a carbohydrate-electrolyte drink on the restoration of fluid balance and exercise capacity in a hot, humid environment. *Eur J Appl Physiol.* 2008; 104: 633-642.
4. Desbrow B et al. Comparing the rehydration potential of different milk-based drinks to a carbohydrate–electrolyte beverage. *Appl Physiol Nutr Metab.* 2014; 39: 1366-1372

5. Seery S, Jakeman P. A metered intake of milk following exercise and thermal dehydration restores whole-body net fluid balance better than a carbohydrate-electrolyte solution or water in healthy young men. *Br J Nutr.* 2016; 116:1013-1021.
6. Volterman KA et al. Effect of milk consumption on rehydration in youth following exercise in the heat. *Appl Physiol Nutr Metab.* 2014; 39: 1257-1264.
7. James LJ et al. Effect of milk protein addition to a carbohydrate-electrolyte rehydration solution consumed after exercise in the heat. *Br J Nutr* 2011; 105, 393–399.
8. Volterman KA et al. The effect of post exercise milk protein intake on rehydration of children. *Pediatr Exerc Sci.* 2016; 28: 286-295.
9. Maughan RJ et al. A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index. *Am J Clin Nutr.* 2016; 103: 717-723.
10. Elliot TA et al. Milk ingestion stimulates net muscle protein synthesis following resistance exercise. *Med Sci Sports Exerc.* 2006; 38: 667-674.
11. Hartman JW et al. Consumption of fat-free fluid milk following resistance exercise promotes greater lean mass accretion than soy or carbohydrate consumption in young novice male weightlifters. *Am J Clin Nutr.* 2007; 86: 373-381.
12. Wilkinson SB et al. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am J Clin Nutr.* 2007; 85:1031-1040.
13. Josse AR et al. Body composition and strength changes in women with milk and resistance exercise. *Med Sci Sports Exerc* 2010; 42: 1122-1130.
14. Hayes A & Cribb PJ. Effect of whey protein isolate on strength, body composition and muscle hypertrophy during resistance training. *Curr Opin Clin Nutr Metab Care.* 2008; 11: 40-44.
15. Cockburn E et al. Acute milk-based protein-CHO supplementation attenuates exercise-induced muscle damage. *Appl Physiol Nutr Metab.* 2008; 33: 775-783.
16. Cockburn E et al. Effect of milk-based carbohydrate-protein supplement timing on the attenuation of exercise-induced muscle damage. *Appl Physiol Nutr Metab.* 2010; 35: 270-277.
17. Cockburn E et al. Effect of volume of milk consumed on the attenuation of exercise-induced muscle damage. *Eur J Appl Physiol.* 2012; 112: 3187-3194.
18. Rankin P et al. The effect of milk on the attenuation of exercise-induced muscle damage in males and females. *Eur J Appl Physiol.* 2015; 115: 1245-1261.
19. Lunn WR et al. Chocolate milk and endurance exercise recovery: protein balance, glycogen and performance. *Med Sci Sports Exerc.* 2012; 44: 682-691.
20. Barry DW et al. Acute calcium ingestion attenuates exercise-induced disruption of calcium homeostasis. *Med Sci Sports Exerc.* 2011; 43: 617-623.
21. Haakonssen EC et al. The effects of a calcium-rich pre-exercise meal on biomarkers of calcium homeostasis in competitive.



by



**European
Milk
Forum**

www.milknutritiousbynature.eu